



RECEIVED
OCT 09 2001
TECH CENTER 1600/2900

Version with markings to show changes made

COMPOUNDS FOR DEACTIVATING PHOSPHOLAMBAN FUNCTION ON CA-ATPASE (PHOSPHOLAMBAN INHIBITORS)

This application is a divisional of Appl. No. 09/252,063, filed February 18, 1999, presently pending, which is a continuation-in-part of U.S. Appl. No. [08/937,119] 08/937,117, filed on September 24, 1997, now abandoned. The entirety of each of these applications is incorporated by reference herein.

Field of the invention

The present invention relates to determining the three-dimensional structure of phospholamban (PLB) using NMR data of sufficiently high resolution for three dimensional structure determination. The invention also relates to methods for rational drug design enabling the design of compounds for deactivating phospholamban based on using the three-dimensional structure data provided on computer readable media, as analyzed on a computer system having suitable computer algorithms. The invention also relates to phospholamban deactivating compounds with structural, physicochemical and spacial characteristics that allow for the interaction of said compounds with specific residues of phospholamban. The interaction prevents the inhibiting effect of phospholamban on Ca-ATPase, making these compounds useful for treating diseases where the Ca-pumping activity of the Ca-ATPase may be diminished, such as congestive heart disease.

Background of the invention

Phospholamban (PLB) is a low molecular weight protein (52 amino acids), present in cardiac, slow-twitch and smooth muscle, which can be phosphorylated by both cAMP- and Ca²⁺/calmodulin-dependent phosphokinases. The amino acid sequences of phospholamban from different species are shown in Figure 1. The phosphorylation/ dephosphorylation of phospholamban has been shown to regulate the Ca²⁺-ATPase of the sarco/endoplasmic reticulum in myocytes (SERCA₂). It has been

shown that phospholamban, in its non-phosphorylated form, binds to a specific region of the large loop in the cytoplasmic domain of SERCA₂ and inhibits this pump by lowering its affinity for Ca²⁺, while the phosphorylated form does not inhibit SERCA₂.

It has been proposed that a region essential for functional association of phospholamban with Ca²⁺-ATPase lies in the cytoplasmic domain of phospholamban, while the transmembrane region anchors PLB to the sarcoplasmic membrane.

During the last decade, efforts have been made to elucidate, at least partially, the secondary structure of PLB either by means of cross-linking experiments or by reconstitution of SERCA₂ with point-mutated PLB, or, finally, by obtaining direct structural information by circular dichroism, laser light scattering photometry - FTIR spectroscopy and NMR spectroscopy). Molecular modelling has been used to formulate hypotheses on the quaternary structure of the transmembrane region in the PLB pentamer. The structural information obtained has been recently reviewed (Arkin, I. T. et al. (1997) *Annu. Rev. Biophys. Biomol. Struct.*, 26, 157-179).

Since PLB i) is an amphiphatic oligopeptide, ii) contains three cysteins, and iii) is prone to pentamerization also *in vitro*, it is not straightforward to find good conditions to study its structure and, in particular, an appropriate solvent system which prevents unspecific aggregation. Therefore, until now NMR studies have been carried out either on short PLB fragments or in organic solvents. In no cases has evidence of a tertiary structure for the cytosolic domain of PLB been found.

Inhibition of CaATPases may play a causative role in cardiac disorders where the calcium levels of myocytes are high. As phospholamban inhibits SR CaATPase this inhibition may be harmful in such disorders. A compound capable of relieving the inhibitory effects of phospholamban on cardiac SR Ca²⁺-ATPase, e.g. by interrupting phospholamban-Ca²⁺-ATPase interaction, would be potentially useful in the treatment of such disorders. There have been very few examples on compounds which can prevent the inhibition of CaATPase by phospholamban in the literature. Such compounds include anti-phospholamban antibodies, some large polyanionic oligopeptides and tannins. No small molecules with specific interactions with phospholamban has been reported.

In the present invention it has been found that phospholamban can assume a well characterized conformation in which it can bind a broad series of small compounds with common structural, physicochemical and spacial characteristics that allow an interaction of the said compounds with specific residues of phospholamban in the defined conformation. This interaction deactivates phospholamban and prevents its inhibiting

effect on Ca-ATPase. The phospholamban deactivating compounds are potentially useful in the treatment of cardiac disorders, where the activation of the SR CaATPase is beneficial.

SUMMARY OF THE INVENTION

The present invention is based on our complete resolution of the three-dimensional structure of the entire cytosolic domain of phospholamban and the ligand binding site therein.

In one aspect the present invention provides compounds capable of relieving the inhibitory effects of phospholamban on cardiac SR Ca^{2+} -ATPase, such compounds thus acting as phospholamban deactivators through direct binding to the phospholamban protein. These compounds have common structural, physicochemical and spacial characteristics that allow for the interaction of said compounds with specific residues of the ligand binding site of phospholamban.

In another aspect the present invention provides a method of deactivating phospholamban which comprises administering to a mammal in need thereof a compound of the invention, as well as a pharmaceutical preparation comprising a compound of the invention together with a pharmaceutically acceptable carrier.

In another aspect, the present invention provides methods for rational drug design enabling the design of phospholamban deactivators based on using the three-dimensional structure data of phospholamban cytosolic domain provided on computer readable media, as analyzed on a computer system having suitable computer algorithms.

In still another aspect, the present invention provides the three-dimensional structure of phospholamban cytosolic domain provided on computer readable media.

Other aspects of the present invention will be apparent to one of ordinary skill in the art from the following detailed description and examples relating to the present invention.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the amino acid sequences of phospholamban from different species (human (SEQ ID NO: 1), pig (SEQ ID NO: 2), dog (SEQ ID NO: 3), rabbit (SEQ ID NO: 4), rat (SEQ ID NO: 5), mouse (SEQ ID NO: 6), chicken (SEQ ID NO: 7)).

Fig. 2 is an illustration of the NMR structure of the cyclic peptide cP226 (SEQ ID NO: 10).

Fig. 3 is an illustration of the NMR structure of PLB(1-36) (SEQ ID NO: 9).

Fig. 4 is an illustration of the model structure of the complex between PLB(1-36) (SEQ ID NO: 9) and cyclic peptide cP226 (SEQ ID NO: 10).

Fig. 5 is an illustration of the main interactions of cP226 (SEQ ID NO: 10) with PLB(1-36) (SEQ ID NO: 9) in the model of the binary complex. The distances between the heavy atoms capable for electrostatic binding, H-bonding, or hydrophobic interactions are shown.

Fig. 6 is an illustration showing the PLB amino acid side chains which form the binding site for ligands, divided to four interactions sites denoted as S1-S4.

Fig. 7 is the summary of the observed sequential and medium-range NOE connectivities for PLB(1-36) (SEQ ID NO: 9).

Fig. 8 is an illustration of the superposition of compound of Example 1c onto the PLB structure.

Fig. 9 shows the quality of the structure of PLB(1-36) (SEQ ID NO: 9) obtained by NOE data. RMSD per residue and the number of restraints per residue is shown.

Fig. 10 is the summary of the observed sequential and medium-range NOE connectivities for cP226 (SEQ ID NO: 10).

Fig. 11 is an illustration of the family of 12 structures of cP226 (SEQ ID NO: 10) deduced from NMR data.

Fig. 12A and 12B show the effect of the compound of Example 1c (50 and 100 μ M) on the Ca^{2+} uptake rate into the cardiac (A) and fast skeletal muscle (B) SR vesicles.

DETAILED DESCRIPTION OF THE INVENTION

Structure of phospholamban (1-36) (SEQ ID NO:9)

The present invention is based on our complete resolution of the three-dimensional structure of the entire cytosolic domain of phospholamban (PLB) and the ligand binding site therein. It was possible to determine the structure of the phospholamban cytosolic domain using a method of NMR spectroscopy wherein the NMR data is of sufficiently high resolution for the three-dimensional structure determination. The method comprises providing the 1-36 a.a. fragment of phospholamban, which comprises the cytosolic domain and 6 amino acids of the transmembrane domain, for the NMR analysis in aqueous solution containing 30% trifluoroethanol. The three-dimensional structure can then be determined from the NMR


data by distance geometry followed by simulated annealing. The method is described in detail in EXAMPLE 1.

It was found that phospholamban (1-36) (SEQ ID NO:9) fragment assumes a conformation characteristic of a helix-turn-helix motif. The residues of the turn are Ile18, Glu19, Met20, and Pro21, which are adjacent to the two phosphorylation sites Ser16 and Thr17. The proline is in a *trans* conformation. Both helices have predominantly charged and polar residues on one side, whereas the other is lipophilic. The hydrophilic side of the N-terminal helix faces always the lipophilic side of the C-terminal helix defining a pocket which could be described as an [amfiphatic] amphipathic armpit. This may mean that in order to interact with SERCA₂, PLB should assume a prolonged position (i.e. the axes of two α -helices should be nearly parallel), while in the bent conformation those charges would not be exposed to the ATPase but eventually to the surface charges of the phospholipid bilayer. The loose relative positioning of the two helices around the mobile central hinge domain is thus a functional feature of PLB. This flexibility may explain also why, in organic solvent, PLB can assume a prolonged structure.

The structure also reveal that the pocket between the hydrophilic side of the N-terminal helix and the lipophilic side of the C-terminal (defined as an [amfiphatic] amphipathic armpit) is an ideal target for small [amfiphatic] amphipathic drug molecules designed with the purpose of deactivating PLB by stabilizing its bent conformation. Such molecules would relieve the inhibitory effect of phospholamban on cardiac SR Ca²⁺-ATPase, and therefore act as a PLB deactivator through direct binding to the active site of PLB.

Structure of cP226 (SEQ ID NO: 10)

In order to find a lead molecule which would interact with PLB a series of peptides were screened. It was found that a cyclic peptide of formula (pI) (SEQ ID NO: 8) was able to bind to PLB and activate the calcium intake in liposomes containing both SERCA₂ and PLB while being inactive in liposomes lacking PLB. It was concluded that the cyclic peptide of formula (pI) binds to unphosphorylated PLB and prevents the inhibition exerted by PLB on SERCA₂ thus acting as a PLB deactivator. The cyclic peptide (pI) has the structure:



X is preferably Tyr or Ala.

The cyclic peptide of formula (pI) wherein X is Tyr was named cP226 (SEQ ID NO: 10). In order to determine the ligand binding site of PLB, the tertiary structure of the cP226 (SEQ ID NO: 10) was resolved by NMR spectroscopy. The method used is described in detail in EXAMPLE 2. The three-dimensional structure of cP226 (SEQ ID NO: 10) show bend-coil-bend motif. The lipophilic side chains of Trp-3, Leu-5, Trp-7 and Leu-8 are clustered on one side of the cyclic peptide, leaving the most of the polar carbonyl and amine groups of the backbone on the other side. The three dimensional coordinates of the cyclic peptide cP226 (SEQ ID NO: 10) are provided in Table I annexed to the present application.

Structure of the complex cP226•PLB(1-36)

On the basis of the resolved tertiary structures of PLB(1-36) (SEQ ID NO: 9) and its ligand cP226 (SEQ ID NO: 10) it was possible to prepare a model of the complex cP226•PLB(1-36) by molecular modelling. The three-dimensional model describes the interactions between PLB(1-36) (SEQ ID NO: 9) and its ligand which are important in binding of ligands to the cytosolic domain of PLB.

The NMR-solved structures of PLB(1-36) (SEQ ID NO: 9) and cP226 (SEQ ID NO: 10) were used as templates for the building of the complex. cP226 (SEQ ID NO: 10) was docked interactively with help of molecular graphics and guided by possible interactions between the two peptides. The structure of cP226 (SEQ ID NO: 10) shows that the peptide has two negative side chains (Glu-4 and Glu-6) on one side while the other side is hydrophobic (Trp-7, Leu-8, Pro-9) (Figure 2). In PLB(1-36) (SEQ ID NO: 9) there is a cluster of three positive side chains (Arg-9, Arg-13, Arg-14) opposed by a mainly hydrophobic surface of the C-terminal helix (e.g. Leu-28, Leu-31, Phe-32, Phe-35) (Figure 3). cP226 (SEQ ID NO: 10) was manually docked onto PLB(1-36) (SEQ ID NO: 9) so that Glu-4 and Glu-6 come to contact with Arg-9, Arg-13 and Arg-14, while at the same time Trp-7, Leu-8 and Pro-9 are near the hydrophobic surface of

PLB C-terminal helix. This gave the starting point for an energy refinement of the complex.

The energy of the obtained complex was minimized by InsightII using the general valence force field (gvff93). Rough minimization was made by the steepest descents method, followed by conjugate gradients and the Newton method.

The structure of the energy minimized complex cP226•PLB(1-36) is shown in Figure 4. The final total energy of the complex was 113 kcal/mol (non-bond dispersion energy -1574 kcal/mol, coulomb energy -690 kcal/mol).

A schematic sketch of the binding mode of cP226 (SEQ ID NO: 10) to PLB is shown in Figure 5. The binding can be described by four binding sites (S1-S4) which bind Glu-4, Glu-6, Trp-7 and Pro-9, respectively (Table II, Figure 6). Glu-4 has electrostatic/H-bonding interactions with Tyr-6, Arg-9 and Arg-13 (S1), Glu-6 binds to Arg-14 (S2), Trp-7 is buried in a hydrophobic pocket (S3) formed mainly by Met-20, Lys-27 and Leu-28 and Pro-9 binds to a hydrophobic cleft (S4) formed mainly by Phe-32 and Phe-35. Besides, Leu-5 is lined by the hydrophobic part of the side chain of Arg-13 and NH of the indole of Trp-7 can form an H-bond to the carbonyl of Arg-13. The positive N-terminal amino group (NH₃⁺) is near the hydrophobic binding site S4.

Table II: Binding of cP226 (SEQ ID NO: 10) to PLB

Site	PLB	cP226 (<u>SEQ ID NO: 10</u>)
S1	Tyr-6, Arg-9, Arg-13	Glu-4
S2	Arg-14	Glu-6
S3	Met-20, Lys-27, Leu-28	Trp-7
S4	Phe-32, Phe-35	Pro-9

Thus, the term "binding site S1" is defined as the space surrounded by amino acid residues Tyr-6, Arg-9 and/or Arg-13, particularly -OH group of Tyr-6, guanidinium group of Arg-9 and/or guanidinium group of Arg-13.

The term "binding site S2" is defined as the space surrounded by amino acid residue Arg-14, particularly guanidinium group of Arg-14.

The term "binding site S3" is defined as the space surrounded by amino acid residues Met-20, Lys-27 and/or Leu 28, particularly the side chains thereof.

The term "binding site S4" is defined as the space surrounded by amino acid residues Phe-32 and/or Phe-35, particularly the phenyl groups thereof.

The three dimensional atom coordinates of phospholamban (1-36) (SEQ ID NO:9) in a conformation which allows binding of cyclic peptide cP226 (SEQ ID NO: 10) are disclosed in Table III annexed to the present application. The three dimensional atom coordinates of the complex between phospholamban (1-36) (SEQ ID NO:9) and the cyclic peptide cP226 (SEQ ID NO: 10) are disclosed in Table IV annexed to the present application. Figure 8 is an illustration of the superposition of one compound of the invention (compound of Example 1c) onto the PLB structure.

Rational drug design

Structure determination methods are also provided by the present invention for rational drug design (RDD) of PLB ligands. Such drug design uses computer modeling programs that calculate different molecules expected to interact with the determined binding sites or other structural or functional domains of PLB. These molecules can then be produced and screened for activity in deactivating PLB according to methods of the present invention.

The present invention reveals the ligand binding site of PLB cytosolic domain heretofore unknown and comprises a distinct three dimensional arrangement of atoms. The atom coordinates of PLB (1-36) (SEQ ID NO: 9) in a conformation which allows binding of a PLB deactivator of the invention to the PLB cytosolic domain are disclosed in Table III. This structure for the first time enables the structure-based design of highly active PLB deactivators. The structure of PLB cytosolic domain provided herein permits the screening of known molecules or designing of new molecules which bind to the ligand binding site of PLB cytosolic domain, via the use of computerized evaluation systems. For example, computer modeling systems are available in which the atomic coordinates of PLB cytosolic domain and the ligand binding site thereof as provided in Table III can be used as input. Thus, a computer readable medium may be encoded with data representing the coordinates of Table III in this process.

The present invention provides a method for identifying a PLB deactivator comprising the steps of:

- i) providing atom coordinates of the structure of PLB cytosolic domain or portion thereof in a computerized modeling system, ii) identifying compounds which are capable of interacting with said structure, and iii) testing the compounds identified or

analogs derived therefrom for the activation of CaATPase in the presence of phospholamban.

In particular, the present invention provides a method for identifying a PLB deactivator comprising the steps of:

i) providing the atom coordinates of the structure of PLB cytosolic domain or portion thereof in a computerized modeling system, ii) identifying compounds which are capable of interacting with at least three of the binding sites S1, S2, S3 and S4 of the PLB cytosolic domain, and iii) testing the compounds identified or analogs derived therefrom for the activation of CaATPase in the presence of phospholamban.

In particular, the present invention provides a method for identifying a PLB deactivator comprising the steps of:

i) providing atom coordinates of the structure of PLB cytosolic domain or portion thereof in a conformation which allows binding of a PLB deactivator to PLB cytosolic domain, in a computerized modeling system, ii) identifying compounds which are capable of said interaction iii) testing the compounds identified or analogs derived therefrom for the activation of CaATPase in the presence of phospholamban.

In the method of the invention candidate molecules may be obtained by carrying out computer-aided molecular design using the three-dimensional structure of the PLB cytosolic domain particularly when complexed with a PLB inhibitor, and in particular the three-dimensional structure at and/or around the binding sites S1 to S4, and synthesising the molecules so-designed.

In the method of the invention candidate molecules can be tested for their ability to deactivate phospholamban, for example, using assays which are described in detail in EXAMPLE 3 or modification thereof.

The invention also provides a computer readable medium having stored therein atom coordinates of the structure of the PLB cytosolic domain or portion thereof in a conformation which allows binding of a PLB deactivator to the PLB cytosolic domain.

As used herein, "computer readable media" refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as optical discs or CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage

media. A skilled artisan can readily appreciate how any of the presently known computer readable media can be used to create a manufacture comprising computer readable medium having stored therein atom coordinates of the structure of the PLB. The choice of the data storage structure will generally be based on the means chosen to access the stored information. Variety of data processor programs and formats can be used to store the atom coordinate data of the present invention on the computer readable medium.

By providing computer readable media having stored therein atom coordinate data of the structure of the PLB, a skilled artisan can routinely access the atom coordinate data of the structure of the PLB cytosolic domain or portion thereof. Computer algorithms are publicly and commercially available which allow a skilled artisan to access this data provided on a computer readable medium and analyze it for structure determination and/or rational drug design. See, e.g. Biotechnology Software Directory, Mary Ann Liebert Publ., New York (1995).

Structural atom coordinates of the PLB (1-36) (SEQ ID NO: 9) presented in Table III may be stored in a computer readable form on a computer readable storage medium for display as a three-dimensional shape or for other uses involving computer-assisted manipulation of the structural coordinates they define. For example, data defining the three dimensional structure of a PLB cytosolic domain or portions thereof may be stored in a computer readable storage medium, and may be displayed as a graphical three-dimensional representation of the protein structure, typically using a computer capable of reading data from said storage medium and programmed with instructions for creating the representation from such data. The invention thus encompasses a machine, such as a computer, having memory which contains data representing the structural coordinates of the PLB protein of the invention, e.g. coordinates presented in Table III, together additional optional data and instructions for manipulating such data. Such data can be used for a variety of purposes, such as the rational drug design. The invention encompasses the coordinates of Table III as well as any translation or rotation or the like thereof which maintains the internal coordinates, i.e. which maintains their intrinsic, internal relationship. Those skilled in the art will appreciate that the coordinates may be subjected to other transformations including, e.g. molecular mechanics calculations such as dynamic simulation, minimization, etc.

For example, a first set of computer readable data defining the three-dimensional structure of PLB cytosolic domain or a portion thereof is combined with a second set of

computer readable data defining the structure of a candidate molecule using a computer programmed with instructions for evaluating the ability of the candidate molecule to associate PLB cytosolic domain protein and/or the location and/or orientation of such association.

The protein structure encoded by the data may be displayed in a graphical format permitting visual inspection of the structure, as well as visual inspection of the structures association with candidate molecules. Alternatively, more quantitative of computational methods may be used. For example, one method of this invention for evaluating the ability of a candidate molecule to associate with PLB cytosolic domain comprises the steps of i) employing computational means to perform a fitting operation between the candidate molecule and the binding sites of PLB, and ii) analyzing the results of said fitting operation to quantify the association between the candidate molecule and the binding sites of PLB.

One method of this invention provides for selecting from a database of chemical structures a compound capable of binding to PLB cytosolic domain. The method starts with structural coordinates defining the three dimensional structure of PLB cytosolic domain or portion thereof. Binding sites of that three dimensional structure are characterized with respect of the favorability of interactions with one or more functional groups. A database of chemical structures is then searched for candidate compounds containing functional groups disposed for favorable interaction with the PLB based on the prior characterization. Compounds having structures which best fit the points of favorable interaction with three dimensional structure are thus identified.

Computer programs for viewing three dimensional structures or manipulating atom coordinates are available and well known for one skilled in the art.

Phospholamban deactivating compounds

The invention provides phospholamban deactivating compounds being capable of associating with any three of the binding sites S1, S2, S3 and S4 of the PLB cytosolic domain. In particular, the invention provides a phospholamban deactivating compound comprising at least any three of the following:

- (a) a first electronegative moiety being capable of associating with the S1 binding site of the [PL] PLB cytosolic domain when the deactivator is bound thereto said binding site comprising Tyr-6, Arg-9 and/or Arg-13,

- (b) a second electronegative moiety being capable of associating with the S2 binding site of the [PL] PLB cytosolic domain when the deactivator is bound thereto said binding site comprising Arg-14, and
- (c) a first hydrophobic moiety being capable of associating with the S3 binding site of the [PL] PLB cytosolic domain when the deactivator is bound thereto said binding site comprising Met-20, Lys-27 and/or Leu-28.
- (d) a second hydrophobic moiety being capable of associating with the S4 binding site of the [PL] PLB cytosolic domain when the deactivator is bound thereto said binding site comprising Phe-32 and/or Phe-35.

In particular, the invention provides a phospholamban deactivating compound comprising at least any three of the following:

- (a) a first electronegative moiety being capable of forming a hydrogen bond with the -OH group of Tyr-6, a salt bridge with the guanidinium group of Arg-9 and/or a salt bridge with the guanidinium group of Arg-13, of the [PL] PLB cytosolic domain when the deactivator is bound thereto,
- (b) a second electronegative moiety being capable of forming a salt bridge with the guanidinium group of [Arg-13] Arg-14, of the [PL] PLB cytosolic domain when the deactivator is bound thereto,
- (c) a first hydrophobic moiety being capable of associating with a hydrophobic pocket created by Met-20, Lys-27 and/or Leu-28, of the [PL] PLB cytosolic domain when the deactivator is bound thereto and.
- (d) a second hydrophobic moiety being capable of associating with a hydrophobic pocket created by Phe-32 and/or Phe-35, of the [PL] PLB cytosolic domain when the deactivator is bound thereto.

In particular, the invention provides a phospholamban deactivating compound comprising:



- (a) a first electronegative moiety being capable of forming a hydrogen bond with the -OH group of Tyr-6, a salt bridge with the guanidinium group of Arg-9 and/or a salt bridge with the guanidinium group of Arg-13, of the [PL] PLB cytosolic domain when the deactivator is bound thereto,

(b) a second electronegative moiety being capable of forming a salt bridge with the guanidinium group of [Arg-13] Arg-14, of the [PL] PLB cytosolic domain when the deactivator is bound thereto and

(c) a first hydrophobic moiety being capable of associating with a hydrophobic pocket created by Met-20, Lys-27 and/or Leu-28, of the [PL] PLB cytosolic domain when the deactivator is bound thereto and.

Therefore, for a phospholamban deactivator of to the invention, at least three of the first electronegative moiety of (a), the second electronegative moiety of (b), the first hydrophobic moiety of (c) and the second hydrophobic moiety of (d) are capable of forming said hydrogen bond and/or salt bridges of (a), said salt bridge of (b), said associations with the hydrophobic pocket of (c), and said associations with the hydrophobic pocket of (d), respectively, with said groups or hydrophobic pockets of PLB, at the same time.

Phospholamban deactivating compounds of the invention include, but are not limited to, compounds of formula (I) or (II):



in which

R₁ is hydrogen, alkyl, alkenyl, aryl, arylalkyl, hydroxyalkyl, halogenalkyl, alkoxy, COR₁₀, CONR₁₀R₁₁, OR₁₀, S(O)_mR₁₀, NR₁₀COR₁₁ or NR₁₀R₁₁, where R₁₀ is hydrogen, alkyl, alkenyl, aryl, arylalkyl, hydroxyalkyl, halogenalkyl, alkoxy or hydroxy and R₁₁ is hydrogen, alkyl, aryl, arylalkyl, alkoxy, aryloxy, hydroxy or acyl, or in case where X is NR₁₁, can R₁ also be carboxylalkyl,

R₆ is hydrogen, alkyl, alkenyl, aryl, arylalkyl,

R₂ and R₇ mean hydrogen, alkyl, aryl, arylalkyl, alkenyl, COR₁₀, CONR₁₀R₁₁, halogen, trifluoromethyl, nitro or cyano, where R₁₀ and R₁₁ are defined as above,

R₃ is hydrogen, alkyl, aryl or arylalkyl,



A means alkyl or substituted alkyl,

m is 0-2 and n is 1-3,

Y means O, NR₁₁ or S, where R₁₁ is the same as above,

X means O, NR₁₁ or S, where R₁₁ is the same as above,

R₄, R₅, R₈ and R₉ mean independently one of the following groups:




or in case where X is NR₁₁, can R₄, R₅, R₈ and R₉ also independently mean HOOC-, R₁₂OOC-, H₂NCO- or HOHNCO- wherein R₁₂ means alkyl, arylalkyl or aryl, and wherein each aryl residue defined above by itself or as part of another group may be substituted, and pharmaceutically acceptable salts and esters thereof.

The compounds of formula (I) or (II) share the structural features which allow them to associate with the ligand binding site of PLB cytosolic domain thereby relieving the inhibitory effects of PLB on cardiac SR Ca²⁺-ATPase.

Compounds of formula (I) or (II) can be prepared from the 1,3-dihydroxy substituted heteroaromatics by alkylation of the dihydroxy compounds by suitable alkylating agents, for example by chloroacetonitrile or bromoacetic ester according to the following Scheme 1, wherein R_1, R_2, R_3, X and Y are the same as defined above, R' is a protecting group for the hydroxyl, e.g. methyl, benzyl or tetrahydropyranyl.


SCHEME 1



The cyano compound (IV) described above is used to prepare the 1,2,4-oxadiazole and 1,2,4-thiadiazole derivatives using the methods described in Kohara et al. (1996) J. Med. Chem. [1996], 39, 5228-5235.

The syntheses are shown in Scheme 2, wherein R_1 , R_2 , R_3 , X and Y are the same as defined above.

SCHEME 2




The other heterocyclics as groups R_4 , R_5 , R_8 and R_9 are prepared as described in Ferrari, B. et al. (1994) Bioorg. Med. Chem. Lett., [1994,]4, 45-50.


The dihydroxyaromatics (III) are made by use of the literature methods. The coumarins (XIV), (XVI) and (XX) are made by the use of the Knoevenagel condensation or von Pechmann reaction as presented in Scheme 3 and 4, where R_1, R_2

and R₃, are the same as defined above, Z is alkyl, aryl, arylalkyl or alkenyl and R' is a protecting group for the hydroxyls e.g. methyl, benzyl or tetrahydropyranyl.

SCHEME 3:








SCHEME 4.




The quinolinones are prepared by the Knorr reaction as described in Scheme 5, wherein R_1, R_{11} and R_3 are the same as defined above, X is a halogen.

SCHEME 5.



The cyclic compounds (II) can be prepared correspondingly from compound (XXXI) which can be prepared according to the Scheme 6, wherein R₂ and R₆ are the same as defined above, R' is a protecting group for the hydroxyls e.g. methyl, benzyl or tetrahydropyranyl.

SCHEME 6.



Cyclic quinolinone compounds (II) can be prepared correspondingly from (XXVI) using Scheme 5.

Salts and esters of the compounds, when applicable, may be prepared by known methods. Physiologically acceptable salts are useful as active medicaments, however, preferred are the salts with alkali or alkaline earth metals. Physiologically acceptable esters are also useful as active medicaments. Examples are the esters with aliphatic or aromatic alcohols.

The term "alkyl" as employed herein by itself or as part of another group includes both straight and branched chain radicals of up to 18 carbon atoms, preferably 1 to 8 carbon atoms, most preferably 1 to 4 carbon atoms. The term "lower alkyl" as employed herein by itself or as part of another group includes both straight and branched chain radicals of 1 to 7, preferably 1 to 4, most preferably 1 or 2 carbon atoms. Specific examples for the alkyl and lower alkyl residues, respectively, are methyl, ethyl, propyl, isopropyl, butyl, tert. butyl, pentyl, hexyl, octyl, decyl and dodecyl including the various branched chain isomers thereof.

The term "acyl" as employed herein by itself or as part of another group refers to an alkylcarbonyl or alkenylcarbonyl group, the alkyl and alkenyl groups being defined above.

The term "aryl" as used herein by itself or as part of another group refers to a monocyclic or bicyclic group containing from 6 to 10 carbon atoms in the ring portion. Specific examples for aryl groups are phenyl, naphthyl and the like. "Aroyl" means in a corresponding way an arylcarbonyl group.

The term "alkoxy" as employed herein by itself or as part of another group includes an alkyl group as defined above linked to an oxygen atom. "Aryloxy" means in a corresponding way an aryl group linked to an oxygen atom.

The term "substituted" as used herein in connection with various residues refers to halogen substituents, such as fluorine, chlorine, bromine, iodine or trifluoromethyl group, amino, alkyl, alkoxy, aryl, alkyl-aryl, halogen-aryl, cycloalkyl, alkylcycloalkyl,

hydroxy, alkylamino, alkanoylamino, arylcarbonylamino, nitro, cyano, thiol, or alkylthio substituents.

The "substituted" groups may contain 1 to 3, preferably 1 or 2, most preferably 1 of the above mentioned substituents.

Compound of the invention may be administered to a patient in therapeutically effective amounts which range usually from about 0.1 to 500 mg per day depending on the age, weight, condition of the patient, administration route and the phospholamban deactivator used. The compounds of the invention can be formulated into dosage forms using the principles known in the art. It can be given to a patient as such or in combination with suitable pharmaceutical excipients in the form of tablets, dragees, capsules, suppositories, emulsions, suspensions or solutions. Choosing suitable ingredients for the composition is a routine for those of ordinary skill in the art. It is evident that suitable carriers, solvents, gel forming ingredients, dispersion forming ingredients, antioxidants, colours, sweeteners, wetting compounds and other ingredients normally used in this field of technology may be also used. The compositions containing the active compound can be given enterally or parenterally, the oral route being the preferred way. The contents of the active compound in the composition is from about 0.5 to 100 %, preferably from about 0.5 to about 20 %, per weight of the total composition.

The following Examples serve purely as an illustration of the various aspects of this invention, and is not intended to be limiting the present invention.

EXAMPLE 1. Structure Of The Cytosolic Domain of Phospholamban

Phospholamban (1-36) (SEQ ID NO: 9) synthesis, purification and characterization

The cytosolic part of phospholamban peptide having the amino acid sequence MEKVQYLTRSAIRRASTIEMPQQARQKLQNLFINFC (SEQ IN NO: 9) was synthesized with an automated peptide synthesizer (Perkin-Elmer, Applied Biosystems 431A) using the fluorenylmethoxycarbonyl strategy. The synthesis was started from the hydrophobic C-terminal end. The side chain protecting groups employed during the synthesis were: Trityl (Trt) for Asn, Gln, and Cys, tert-Butoxy (OtBu) for Glu, tert-Butyl (tBu) for Ser, Thr and Tyr, tert-Butoxycarbonyl (Boc) for Lys and 2,2,5,7,8-Pentamethyl chroman-6-sulphonyl (Pmc) for Arg.

The amount of the preloaded resin was 100 μ mol and the quantity of the amino acids at each step of the synthesis was 1 mmol. This is a 10 times excess as compared to the amount of the resin loaded.

The peptide cleavage from the resin support (originally preloaded Wang resin) was performed in methylene chloride containing 5 % TFA, 0.2 % β -mercaptoethanol, 0.2 % thio anisole and 0.2 % dimethylsulfide. Cleavage of the side chain protection of the peptide was carried out in a mixture of ethanedithiol : thioanisole : water : trifluoroacetic acid = 250 μ l : 500 μ l : 500 μ l : 10 ml for 1.5 hours at room temperature. After that the peptide was precipitated and washed three times with diethylether, and lyophilized.

The synthesized crude phospholamban (1-36) (SEQ IN NO: 9) peptide was prepurified by high performance liquid chromatography (HPLC) technique with an analytical reversed phase (RP) column (C₈, 20 μ m, 4.6 mm x 30 mm i.d. Perkin-Elmer, Applied Biosystems Brownlee TM column). A linear gradient of acetonitrile (0-100% in 30 min.) in 0.1% TFA was used for elution.

The repurified peptide was further purified by HPLC RP-chromatography using a C₁₈ Kromasil, 5 μ m (1.0 x 25 cm) column. The peptide was eluted using a stepwise gradient of acetonitrile, 0.075% TFA (3-30%, in 10 min, 30-50% in 120min.) in 0.1% TFA.

The purified phospholamban (1-36) peptide (SEQ IN NO: 9) was characterized by SDS-PAGE followed by Coomassie brilliant blue staining. Western blot analysis was done by using the commercial monoclonal anti-PLB antibody (Upstate Biotechnology). The purified RP-chromatography peaks containing the peptide was further analyzed by mass spectrometry (MALDI-TOF) in reflector mode with a BIFLEX™ mass spectrometer using a 337nm nitrogen laser. The samples were applied in a solution containing 30% acetonitrile/0.1% TFA together with a droplet of sinapinic acid matrix for mass spectrometry analysis. The total amount of purified protein was estimated according to Bradford and also based on RP-chromatography using β -lactoglobulin as a standard reference. The purified peptides were lyophilized and the dry powder was estimated by weight before analysis of the 3-dimensional structure.

Obtaining NMR spectra of phospholamban (1-36) (SEQ IN NO: 9)

^1H -NMR spectra were acquired at 400.13 MHz and at 599.86 MHz on a Bruker ARX400 and a Varian UNITY 600 NMR spectrometer respectively. 1D and 2D NMR spectra were obtained for a 3 mM solution of the 36-a.a. fragment of PLB in the solvent mixture $\text{H}_2\text{O}:\text{D}_2\text{O}:\text{d}_3\text{-TFE}$ (63:7:30) containing 6 mM d_{10} -DTT to prevent disulphide formation. The pH was adjusted to 3.00 ± 0.02 (uncorrected for deuterium isotope effects) with microliter amounts of NaOD. COSY, TOCSY (30 - 90 ms) and NOESY (40 - 400 ms) spectra were recorded at 2, 7, 17 and 27°C, by the States-TPPI method using a spectral width of 8.5 ppm. The 2D data was weighted and Fourier transformed to 2 k x 1 k real point matrices. The transmitter presaturated (2.0 s) residual solvent line was reduced by deconvolution. The spectra were referenced to the residual solvent signal (4.75 ppm at 27°C, -10 ppb/°C). A series of ten 1D spectra was acquired at different temperatures (ranging from 2 to 47°C).

Assignment of the NMR spectra of phospholamban (1-36) (SEQ ID NO:9)

The spin-system and sequential assignments were derived according to Wüthrich, K. et al. (1986) *NMR of Proteins and Nucleic Acids*, John Wiley & Sons, Inc., New York, by use of COSY, TOCSY and NOESY spectra acquired at 12, 17 and 27°C. Differences in the temperature dependences of the amido proton chemical shifts were sufficient to unravel resonance overlap. Stereospecific assignments for non-degenerated methylenes were deduced from coupling constants $J_{\text{H}\alpha\text{H}\beta}$ measured from the COSY spectra and from intra residual NOE-cross peak intensities.

Phospholamban (1-36) (SEQ ID NO:9) structure generation and refinement

A series of NOESY spectra was acquired at 17°C with five different mixing times (50, 80, 120, 160, and 200 ms). The integrated cross peak intensities (I) were used in a NOESY-built-up-analysis. Distance restraints were extracted from the initial slope of a second-order polynomial curve fitted to the volumes of the cross peaks integrated from the NOE-series, with the initial condition $I_{(\tau_m=0)}=0$. Intra methylene and sequential NOEs served for the calibration. The distances were initially classified as short (1.8-2.5 Å), medium (1.8-3.5 Å) or long (3.0-6.0 Å) for the generation of the first set of structures. When a distance could not be calculated from the built-up curve, owing to a partial (> 20 %) overlap, a poor signal-to-noise ratio or disturbances, it was only required that the distance was < 5.0 Å. The upper bounds were extended by 1.0 Å for each pseudo atom. The restraint data were supplemented with distance restraints, which

were based on strong, medium and weak NOEs, from the 150 ms NOE-spectrum acquired at 12°C.

Coupling constants (J) were measured by the J -doubling method (McIntyre, L. et al. (1992) *J Magn Reson* 96, 425-431) from fine structures of COSY cross peaks. Dihedrals ϕ and χ , which were characterised by intermediate J , were not constrained but small and large $J_{\text{NH}\alpha}$ and $J_{\text{H}\alpha\text{H}\beta}$ were related to staggered conformers (± 30 degrees) on the basis of Karplus functions and intra residual NOEs (Karplus, M. (1963) *J. Am. Chem. Soc.*, 85, 2870). The H-H distance and dihedral angle restraints were calculated and the data were imported into the software InsightII (Molecular Simulations, Inc.) in order to generate, evaluate and refine the structures. Simulated NOESY spectra were back calculated. The protein coordinate files were analysed by the software PROMOTIF v2.0 (Hutchinson, G. (1995), v2.0 Ed., available by anonymous ftp on 128.40.46.11).

Structures were generated by distance geometry (DGII) followed by simulated annealing (force field AMBER) (Havel, T. et al. (1979) *Biopolymers* 18, 73). A set of structures was computed. The structures with the least restraint violations were used to back calculate NOE-matrices. If the $\text{H}\alpha$ -chemical shift of consequent residues in the segment characterised by NOEs typical of α -helices departed from the corresponding random coil value more than -0.2 ppm, also the correspondent dihedrals ψ were constrained (± 60 degrees). A new set of structures was subsequently calculated. From this new family of structures only those structures with no violations over 0.2 Å were accepted.

Assignment results

The complete spin-system and sequential assignments were obtained under the experimental conditions described. The assignments are listed in Table V (annexed to the present application) showing H-chemical shifts of PLB (1-36) (SEQ ID NO:9) in 54% H_2O / 6% D_2O / 30% d_3 -TFE, pH 3.05 at 17°C, wherein the staggered conformations are denoted by a line under the chemical shift of C_βH in an anti-configuration and by a dashed line in a gauche-configuration (-60 degree) to C_αH .

Fig. 7 is a summary of the observed sequential and medium-range NOE connectivities for PLB(1-36) (SEQ ID NO: 9) in 54% H_2O / 6% D_2O / 30% d_3 -TFE, at pH 3.05 at 17°C., wherein assignments were made from NOESY spectra acquired at 120, 160 and 200 ms mixing times. Sequential NOEs are represented by shaded blocks. Medium-range NOEs are represented by arrows connecting the appropriate residues.

Open circles denote $^3J_{\text{NH}\alpha\text{CH}}$ coupling constants smaller than 7 Hz. The secondary shift ($\Delta\delta$) of αCH is defined as the difference between the observed chemical shift and the random coil chemical shift for each residue. Negative (upfield) $\Delta\delta$ values are associated with α -helical secondary structure and positive (downfield) $\Delta\delta$ values with β -structure according to Wishart et al., *Biochemistry* (1992), 31, 1647-1651.

In total 723 NOEs were assigned. All the 34 possible intra $\text{NH}-\text{C}\alpha\text{H}$ correlations were observed in the finger print region. Most of the corresponding NOEs were fairly strong and comparable to sequential $\text{NH}_{i+1}-\text{C}\alpha\text{H}_i$ NOEs (Fig. 7). Many sequential $\text{NH}_i-\text{NH}_{i+1}$ and $\text{NH}_i-\text{NH}_{i+2}$ NOEs were present. Numerous $\text{C}\alpha\text{H}_i-\text{NH}_{i+3}$ and some $\text{C}\alpha\text{H}_i-\text{NH}_{i+4}$ NOEs were crowded in the finger print region of the NOESY spectra. Furthermore, there were a number of $\text{C}\alpha\text{H}_i-\text{C}\beta\text{H}_{i+3}$ cross peaks. NOEs derived from interactions longer than $i > i+4$ were observed only for protons of Met20.

J -couplings between NH and $\text{C}\alpha\text{H}$ were small for most residues. Due to overlap of resonances or weak intensity of COSY cross peaks it was not possible to measure accurate values for all residues but the couplings were below 7-8 Hz with the exception of the residues at the N- and C-terminus and in the center of PLB (1-36) (SEQ ID NO: 9).

The central region of the PLB(1-36) (SEQ ID NO: 9) does not show a helical character. Namely, the $\delta_{\text{C}\alpha\text{H}}$ values of the residues Glu19, Met20 and Pro21 were not significantly smaller than their random coil values. Glu19 and Met20 were mostly devoid of the NOEs typical of a helical structure, and there were unambiguous strong sequential NOEs between $\text{C}\alpha\text{H}$ of Glu19 and NH of Met20, and between $\text{C}\alpha\text{H}$ of Met20 and $\text{C}\beta\text{H}$ s of Pro21. Furthermore, the NOE between $\text{C}\alpha\text{H}$ of Ile18 and NH of Glu19 is strong, even when partially buried in the crowded finger print region. All this implies that the central region of PLB(1-36) (SEQ ID NO: 9) assumes an extended-like conformation. The extended segment is, nevertheless, short. Thr17 and Gln22 show NOEs and coupling constants characteristic of residues in an α -helix, and there are NOEs from the side chain protons of Glu19 and Met20 to the protons of the adjacent residues in the N- and C-terminal helices. We conclude that the N- and C-terminal α -helices are separated by a turn at Ile18, Glu19, Met20, and Pro21. The proline is in a *trans*-conformation. A tight turn, which would result the axes of the N- and C-terminal helices being parallel, is not possible. There were no unambiguous NOEs between the N- and C-terminal helices.

Structure of PLB (1-36) (SEQ ID NO: 9)

The structure of PLB (1-36) (SEQ ID NO: 9) was determined from 599 distances and 50 dihedral restraints excluding those that were defined more accurately by the covalent structure alone. These redundant NOE-derived restraints were consistent with the covalently imposed distance limits, which indicated that the calibration of distances was reasonable. On average there were 16.6 non-trivial NOE-derived restraints per residue. The residues Lys3-Ile18 of the N-terminal helix had on average a few restraints less per residue than the residues Gln22-Cys36 in the C-terminal helix. This is at least partly due to the fact that there were on average more protons with non-degenerated shifts per residue in the C-terminal helix than in the N-terminal helix (Table V). For the residues Ile18-Pro21, which confine the turn, there were about as many restraints per residue as there were for the residues in the N-terminal helix.

The structure generation resulted in a family of structures all of which show a helix-turn-helix motif. The root mean square deviation was computed from the family of 20 structures with no distance violations above 0.2 Å and no dihedral violations. Fig. 9 shows the quality of the structure of PLB(1-36) (SEQ ID NO: 9) obtained by NOE data wherein RMSD per residue and the number of restraints per residue is shown.

Since no unambiguous long range NOEs were found between the N- and C-terminal helices, the family of structures displayed a dispersion of atomic coordinates in the remote parts of the N- and C-terminal helices. The mutual orientation of the helices was constrained only by the short range distance restraints in the turn. Therefore, RMSD per residue was computed separately i) for the N-terminal helix plus the turn (a.a. 1-21) and ii) for the C-terminal helix plus the turn (18-36). The RMSD represented roughly an inverse correlation with the number of restraints per residue, as expected. In average the atoms in the N-terminal helix were defined to a precision of 1.3 Å (backbone only) and of 2.3 Å (all atoms) and in the C-terminal helix to a precision of 0.8 Å (backbone only) and of 1.9 Å (all atoms). The smallest number of distance violations (below 0.2 Å) were observed for structures in which the segment from C α of Glu19 to C α of Pro21 is extended, the side chain of Met20 sticks out approximately parallel to the C-terminal helix and the side chain of Glu19 points almost in the opposite direction. In these structures, the plane of the peptide bond Ile18-Glu19 is approximately orthogonal to the plane of the extended segment.

Owing to the structural mobility of the turn, the family of structures displays a dispersion in the relative position of the N- and C-terminal helices. The dispersion is, nevertheless, limited. When the different structures of the family are superimposed on the C α of the residues in the C-terminal helices, the axis of the N-terminal is dispersed in a cone with an opening of approximately 90 degrees, and with a relative angle of

about 80 degrees. Similar mutual orientations for two sequential helices, one of which transmembran and the other anfiphatic, have been found or hypothized for many small membrane bound proteins or peptides (Stopar, D. et al. (1996) *Biochemistry*, 35 (48), 5467-5473).

In some of the structures, the side chain ϵ NH of Arg9, Arg13 and Arg14, whose chemical shifts are nearly independent of T , make hydrogen bonds with the adjacent side chain oxygen of Ser10, Ser16 and Thr17. For Arg25, with large $\Delta\delta_{\text{NH}}(T)$, there were no obvious candidates for hydrogen bond donors. The side chain NH_2 of the glutamines and asparagines could form hydrogen bond net works parallel to the helical axis.

With regards to the phosphorylation of PLB, we find important that the phosphorylation site Ser16 is readily accessible and exposed to the solvent. Thr17, on the N-terminal helix, is facing the C-terminal helix and appears less exposed to the solvent than Ser16. Due to the pitch of the α -helix, Arg13 and Arg14 are also exposed with orientations that lag 60 degrees in phase with respect to Ser16 and Thr17 on the same side of the helix. The presence of positively charged residues in the vicinity of a serine or threonine residue is often seen in a substrate for phosphokinases (e.g. in Troponin I).

We find that one side of the N-terminal helix is predominantly polar or hydrophilic. The same is true for the C-terminal helix, whose polar residues are located on one side while leaving the other side dominated by lipophilic side chains. Owing to the relative orientation of the helices, it can be noted that the hydrophilic side of the N-terminal helix faces always the lipophilic side of the C-terminal helix. This defines a pocket which could be described as an [amfiphatic] amphipathic armpit. In this pocket, the relative position of the polar residues on the N-terminal and of the lipophilic residues on the C-terminal can be conveniently described by use of two centroids calculated by averaging the coordinates of selected side chains. The coordinates of the side chains of Arg13, Arg9 and Tyr6 were used to build the centroid relative to the N-terminal and the coordinates of the side chains of Phe32 and Phe35 were used to build up the centroid relative to the C-terminal. The distance between these two centroids was calculated for every structure of the family and was 18.5 ± 4.5 Å. The refined structure of PLB (1-36) (SEQ ID NO: 9) is shown in Figure 3.

EXAMPLE 2. Structure Of The Cyclic Peptide cP226 (SEQ ID NO: 10)

Peptide synthesis, cyclization and purification

The linear peptide having the amino acid sequence CYWELEWLPCA (SEQ ID NO: 10) was synthesized with an automated peptide synthesizer (Perkin-Elmer, Applied Biosystems 431A) using the fluorenylmethoxycarbonyl-strategy. The synthesis was started from the carboxy-terminal end. The side chain protecting groups employed during the synthesis were: Trityl (Trt) for Cys, tert-Butoxy (OtBu) for Glu, and tert-Butyl (tBu) for Tyr.

The amount of the preloaded resin was 100 μ mol and the quantity of the amino-acids at each step of the synthesis was 1 mmol.

The peptide cleavage from the resin support (originally preloaded Wang resin) as well as cleavage of the side chain protection groups was carried out in a mixture of ethanedithiol : thioanisole : water : trifluoroacetic acid = 250 μ l : 500 μ l : 500 μ l : 10 ml for 1.5 hours at room temperature. After that the peptide was precipitated and washed three times with diethylether, and lyophilized.

The amino-acids and the preloaded resins used for the peptide synthesis were obtained from Novabiochem. Trifluoroacetic-acid (TFA) was produced by Perkin-Elmer, ethanedithiol (EDT) and thioanisole were manufactured by Fluka.

The cyclic peptide cP226 (SEQ ID NO: 10) was reconstituted from the linear CYWELEWLPCA peptide (SEQ ID NO: 10) by dissolving 0.5 mg/ml the purified peptide into 10 mM $(\text{NH}_4)_2\text{CO}_3$ and the oxidation of the SH groups of the two cysteine residues to form intramolecular disulphide bridge was achieved by leaving the solution at room temperature for 1-2 days. The reaction was followed by HPLC chromatography from the peaks of the linear and the cyclic peptide was varied as a function of time.

The peptides, both the linear and the cyclic cP226 (SEQ ID NO: 10), were purified and separated by reverse phase HPLC-chromatography (C8, Aquapore Octyl, 30 μ m, 10x100 mm, Perkin-Elmer) using 30 min linear gradient from 0.1 % TFA to 100 % acetonitrile. The obtained peptides were characterized by mass spectrometry.

NMR spectra of cP226 (SEQ ID NO: 10)

^1H -NMR spectra were acquired at 400.13 MHz and at 599.86 MHz on a Bruker ARX400 and a Varian UNITY 600 NMR spectrometer respectively. 1D and 2D NMR spectra were obtained for a 1 mM solution of the cyclic peptide in water. The pH was adjusted to 6.50 ± 0.02 (uncorrected for deuterium isotope effects) with microliter amounts of NaOD. COSY, TOCSY (30 - 90 ms) and NOESY (200 - 400 ms) spectra

were recorded at 5, 10, 15 and 27°C, by the States-TPPI method, using a spectral width of 8.5 ppm. The 2D data was weighted and Fourier transformed to 2 k × 1 k real point matrices. The transmitter presaturated (2.0 s) residual solvent line was reduced by deconvolution. The spectra were referenced to the residual solvent signal (4.75 ppm at 27°C, -10 ppb/°C).

Assignment of the NMR spectra of cP226 (SEQ ID NO: 10)

The spin-system and sequential assignments were derived according to Wüthrich as in EXAMPLE 1, by use of COSY, TOCSY and NOESY spectra acquired at 5, 10 and 27°C. Differences in the temperature dependences of the amido proton chemical shifts were sufficient to unravel resonance overlap. Stereospecific assignments for non-degenerated methylenes were deduced from coupling constants $J_{H\alpha H\beta}$ measured from the COSY spectra and from intra residual NOE-cross peak intensities.

cP226 (SEQ ID NO: 10) Structure Generation And Refinement

A series of NOESY spectra was acquired at 10°C with different mixing times (200, 300, 400 ms). The integrated cross peak intensities (I) were used in a NOESY-built-up-analysis. Distance restraints were extracted from the initial slope of a second-order polynomial curve fitted to the volumes of the cross peaks integrated from the NOE-series, with the initial condition $I(\tau_m=0)=0$. Intra methylene and sequential NOEs served for the calibration. The distances were initially classified as short (1.8-2.5 Å), medium (1.8-3.5 Å) or long (3.0-6.0 Å) for the generation of the first set of structures. When a distance could not be calculated from the built-up curve, owing to a partial (> 20 %) overlap, a poor signal-to-noise ratio or disturbances, it was only required that the distance was < 5.0 Å. The upper bounds were extended by 1.0 Å for each pseudo atom. The restraint data were supplemented with distance restraints, which were based on strong, medium and weak NOEs, from the 300 ms NOE-spectrum acquired at 10°C.

Coupling constants (J) were measured by the J -doubling method from fine structures of COSY cross peaks. Dihedrals ϕ and χ , which were characterised by intermediate J , were not constrained but small and large $J_{NH\alpha}$ and $J_{H\alpha H\beta}$ were related to staggered conformers (± 30 degrees) on the basis of Karplus functions and intra residual NOEs. The H-H distance and dihedral angle restraints were calculated. Finally, the data were imported into the software InsightII (Molecular Simulations,

Inc.) in order to generate, evaluate and refine the structures. Simulated NOESY spectra were back calculated. The protein coordinate files were analysed by the software PROMOTIF v2.0.

Structures were generated by distance geometry (DGII) followed by simulated annealing (force field AMBER). A set of 30 structures was computed. The structures with the least restraint violations were used to back calculate NOE-matrices. Based on the comparison of the back-calculated and experimental NOE-spectra it became possible to unambiguously identify more NOEs and impose corresponding distance restraints. A new set of 30 structures was subsequently calculated. From this new family, 12 structures with at most one violation larger than 0.3 Å were selected and examined further. The distance restraints corresponding to well-resolved cross peaks were refined by an iterative relaxation matrix method (IRMA) based on a structure without restraint violations (>0.2 Å, >0 deg). The upper bounds were kept within at least 10% of the exact distance given by IRMA to take into account the uncertainty in τ_c . The refined restraint set was subsequently used to refine the coordinates by simulated annealing.

The resulting final family of 12 structures was visualized by the graphic software MOLMOL (Koradi, R. et al. (1996) *J Mol Graphics* 14, 51-55).

Assignment results

The complete spin-system and sequential assignments were obtained under the experimental conditions described. It was necessary to run the NOESY experiments at low temperature ($<10^{\circ}\text{C}$) and with relatively long mixing time (>300 ms) to induce enough magnetization transfer and visualize the cross peaks. The assignments are listed in Table VI.

cP226 (SEQ ID NO: 10) spectra displayed chemical shift dispersion over 8.5 ppm. $\text{C}_{\alpha}\text{H}$ shifts ranged from 3.9 to 4.7 ppm and most of the NH shifts were confined between 7.8 and 8.5 ppm, but for Tyr2 and Leu8 the NH resonances were shifted upfield, to 6.9 and 7.3 ppm respectively. There were no signals of methyl groups at very high field (> 0 ppm).

Fig. 10 is a summary of the observed sequential and medium-range NOE connectivities for CP226 (SEQ ID NO: 10) in 90% H_2O / 10% D_2O , pH 6.50 at 10°C . The assignments were made from NOESY spectra acquired at 300 and 400 ms mixing times. Sequential NOEs are represented by shaded blocks. Medium-range NOEs are represented by arrows connecting the appropriate residues. Open circles denote $^3\text{J}_{\text{NH}\alpha\text{CH}}$ coupling constants bigger than 8 Hz. The secondary shift ($\Delta\delta$) of αCH is defined as the difference between the observed chemical shift and the random coil chemical shift for each residue. Negative (upfield) and positive (downfield) $\Delta\delta$ values are associated to the secondary structure according to Wishart et al., *Biochemistry* (1992), 31, 1647-1651.

In total 120 NOEs were assigned. Not all the 9 possible intra $\text{NH}-\text{C}_{\alpha}\text{H}$ correlations were observed in the finger print region. Most of the corresponding NOEs were fairly weak. Only four sequential $\text{NH}_i-\text{NH}_{i+1}$ and one $\text{NH}_i-\text{NH}_{i+2}$ NOEs were present. Some $\text{NH}_{i+n}-\text{C}_{\beta}\text{H}_i$ were visible in the NOESY spectra and facilitated the sequential assignment. The secondary shift ($\Delta\delta$) of αCH , defined as the difference between the observed chemical shift and the random coil chemical shift for each residue, gave evidence of bended structure (Fig. 10). NOEs derived from interactions longer than $i > i+4$ (across the cycle) were observed for protons of the side chains of Trp3 and Leu8 and for protons of Tyr2 and Cys10.

J-couplings between NH and $\text{C}_{\alpha}\text{H}$ (measured from the COSY cross peaks) were large for most residues. The couplings for all residues were above 8 Hz, suggesting that

the ϕ angles have predominantly values of $120 \pm 30^\circ$. The proline was in a trans-conformation.

Structure of cP226 (SEQ ID NO: 10)

The structure of cP226 (SEQ ID NO: 10) was determined from 110 distances and 8 dihedral restraints excluding those that were defined more accurately by the covalent structure alone. These redundant NOE-derived restraints were consistent with the covalently imposed distance limits, which indicated that the calibration of distances was reasonable. On average there were 10 non-trivial NOE-derived restraints per residue. The residues in the central part of the peptide (from Trp3 to Leu8) had more restraint per residue than the average. This is at least partly due to the fact that there were on average more protons with non-degenerated shifts per residue in that zone.

The structure generation resulted in a family of structures all of which show bend-coil-bend motif (MOLMOL). The root mean square deviation was computed from the family of 12 structures with no distance violations above 0.3 Å and no dihedral violations. The small distance restraint violations occurred primarily among the side chain groups, e.g. the tyrosine side chain. This may be a result of excessive mobility in these parts, which could give rise to non-simultaneous NOEs. RMSD per residue was computed and represented roughly an inverse correlation with the number of restraints per residue, as expected. In average the atoms were defined to a precision of 0.96 Å (backbone only) and of 2.02 Å (all atoms) and, if calculated only for the portion from Trp3 to Trp7, to a precision of 0.72 Å (backbone only) and of 1.87 Å (all atoms) (Table VII).

The lipophilic side chains of Trp3, Leu5, Trp7 and Leu8 were clustered on one side of the cyclic peptide, leaving the most of the polar carbonyl and amine groups of the backbone on the other side.

The structure of CP226 (SEQ ID NO: 10) shown in this report was docked on the structure of PLB (1-36) (SEQ ID NO: 9) described above. The comparison of the two structures confirm the hypothesis that the two glutamate residues on CP226 (SEQ ID NO: 10) may participate to the binding by coupling with Arg9 and Arg 13 on PLB (1-36) (SEQ ID NO: 9), exposing the lipophilic cluster of [PC226] CP226 (SEQ ID NO: 10) to the lipophilic outer part of the C-terminal helix of PLB(1-36) (SEQ ID NO: 9).

Fig. 11 shows a family of 12 structures of CP226 (SEQ ID NO: 10) deduced from NMR data. The backbones, the heavy atoms of the Trp3 and Trp7 side chains, the heavy atoms of the Glu4 and Glu6 side chains, the carbon β of Glu4 and Glu6 is shown. The distance between the beta carbons of the two glutamate residues (Fig. 11) was highly conservative ($8.3 \pm 0.9 \text{ \AA}$). This information can be useful to design small molecules in which two acetate residues mimic the position of Glu4 and Glu6 on CP226 (SEQ ID NO: 10). In the same way, the distance of the lipophilic cluster from the two glutamate residues will also be useful in the drug design process.

EXAMPLE 3. Activity assays

Experiment 1. Effect on calcium uptake into the SR vesicles prepared from cardiac and fast skeletal muscle

The inhibitory effect of a given compound on phospholamban can be demonstrated by measuring the effect of the compound on calcium uptake into the SR vesicles prepared from cardiac tissue and into SR vesicles prepared from fast skeletal muscle (psoas m.). Both kind of SR vesicles contain Ca^{2+} -ATPase but the vesicles from the fast skeletal muscle do not contain phospholamban (Hoh JFY, "Muscle fiber types and function", Current Opinion in Rheumatology, 4:801-808, 1992). An increase in the calcium uptake into the SR vesicles prepared from cardiac tissue but not into the SR vesicles prepared from fast skeletal muscle indicates that the compound relieves the inhibitory effect of phospholamban on SR Ca^{2+} -ATPase and thus acts as a phospholamban inhibitor. Since phospholamban represses both the rates of relaxation and contraction in the mammalian heart through its inhibitory effects on the cardiac SR Ca^{2+} -ATPase, a compound relieving these effects is potentially useful in the treatment of heart failure.

Method

Guinea pigs (10-12) were decapited. Their hearts or the psoas muscles were excised, washed in ice-cold 0.9 % NaCl and cut into pieces in a buffer containing 20 mM Tris-maleate, 0.3 M sucrose, pH 7.0. Thereafter tissue pieces were homogenized with Polytron and further with Potter (10 strokes). The homogenate was centrifugated at 1000 g for 15 min at 4 °C. The supernatant was collected and the pellet was resuspended into 5 ml of the buffer (20 mM Tris-maleate, 0.3 M sucrose, pH 7.0) and

recentrifugated at 1000 g for 10 min at 4 °C. The obtained supernatant was combined with the earlier collected supernatant and centrifugated once again at 10 000 g for 20 min at 4 °C. The final supernatant was filtered into a bottle equipped with a magnetic stirrer. KCl was added to the filtered supernatant to achieve the final concentration of 0.6 M (at 4 °C). The obtained solution was centrifugated at 100 000 g for 60 min at 4 °C. The pellet was suspended in 5 ml of the buffer containing 20 mM Tris-maleate, 0.3 M sucrose, pH 7.0 and centrifugated at 100 000 g for 60 min at 4 °C. The obtained pellet was suspended in 5 ml of buffer containing 20 mM Tris-maleate, 0.3 M sucrose, 0.1 M KCl, pH 7.0 and stored at -80 °C until use. The protein concentration was also measured in order to standardise the separately prepared vesicle preparations.

In the calcium uptake assay, the fluorescent indicator, fluo-3 was used to detect the decrease of the extravesicular Ca^{2+} -concentration, when the SR Ca^{2+} ATPase was transferring Ca^{2+} from the extravesicular space into the SR-vesicles.

The SR-vesicles obtained above (50 μg protein/ml) were pre-incubated with or without the test compound at 37 °C for 5 min in the assay buffer containing 40 mM imidazole, 95 mM KCl, 5 mM NaN_3 , 5 mM MgCl_2 , 0.5 mM EGTA, 5 mM potassium oxalate, 2 μM ruthenium red, 5 μM fluo-3, pH 7.0. The free calcium was adjusted to 0.1 μM or to 0.04 μM by CaCl_2 . The reaction was initiated by adding ATP (5 mM). The final reaction volume was 1.5 ml. The fluorescence of reaction mixture was measured for 3 min by using the excitation and emission wavelengths of 510 nm and 530 nm, respectively.

Results

Figures 12A and 12B show the effect of the compound of Example 1c (50 and 100 μM) on the Ca^{2+} uptake rate into the cardiac (A) and fast skeletal muscle (B) SR vesicles. It can be seen that the compound of the invention accelerated the calcium uptake into the cardiac SR vesicles but did not change the calcium uptake into the SR vesicle prepared from the fast skeletal muscle.

Table VIII shows the effects of various compounds of the invention on the Ca^{2+} uptake rate into the cardiac (A) and fast skeletal muscle (B) SR vesicles. The experiments were carried out at 0.1 μM and 0.04 μM free calcium concentrations, respectively.

Table VIII. Stimulation (%) of the Ca^{2+} uptake into the vesicle preparations obtained from the ventricular myocardium (A) and fast skeletal muscle (B) of the guinea-pig heart.

Compound of Example No. (100 μM)	The stimulation (%) of Ca^{2+} uptake	
	A	B
3c**	51	0
2c	26	-1
7c	5	-17
8g*	18	0
11b	28	nd
12	32	nd
13d***	23	nd
14c*	18	nd
18e	13	nd
21	11	nd
23****	20	nd

* 10 μM , ** 20 μM , *** 50 μM , **** 5 μM
nd=not determined

Experiment 2. The effects on the left ventricular pressure derivatives

Method

Guinea-pigs of either sex weighing 300-400 g were used in the study. After the guinea-pig was sacrificed by a blow on the skull and decapitated the heart was rapidly excised. The heart was then rinsed in cold oxygenated perfusion buffer. A cannula was inserted into the aorta and secured with a ligature. Retrograde perfusion began as soon as the heart was placed in a thermostatically controlled moist chamber of the Langendorff apparatus. Modified Tyrode solution (37 °C), equilibrated in a thermostatically controlled bulb oxygenator with carbogen (95 % O_2 and 5% CO_2) was used as a perfusion buffer. The composition of the Tyrode solution was (in mM): NaCl 135; $\text{MgCl}_2 \times 6\text{H}_2\text{O}$ 1; KCl 5; $\text{CaCl}_2 \times 2\text{H}_2\text{O}$ 2; NaHCO_3 15; $\text{Na}_2\text{HPO}_4 \times 2\text{H}_2\text{O}$ 1; glucose 10; pH 7.3-7.4. The experiments were carried out under constant pressure condition (50 mmHg). After a short prestabilization (10 min) a latex balloon (size 4)

was carefully placed into the left ventricle through the left pulmonary vein and the left atrium. The latex balloon was attached to a stainless-steel cannula coupled with a pressure transducer. The latex balloon, the cannula and the chamber of the pressure transducer were filled with ethylene glycol / water (1:1) mixture avoiding any air-bubble. The isovolumetric left ventricular pressure was recorded through the pressure transducer. At the beginning of the experiment, the volume of the balloon was adjusted to obtain a diastolic pressure of approximately 5 mmHg. Before starting the experiment, the heart was allowed to stabilise further for 30 - 50 min with vehicle (0.1% DMSO) in the perfusion buffer.

After 15 min baseline recording various concentrations of the test compound were added to the perfusion buffer at 15 min intervals. The concentration range of 0.3 - 30 μ M was tested. The vehicle concentration (0.1% DMSO) was kept constant throughout the experiment.

Results

The EC₅₀ values and maximum effects (% change from baseline) of various compounds of the invention on left ventricular systolic pressure are given in Table IX.

TABLE IX. The EC₅₀ values and maximum effects (% change from baseline) on left ventricular systolic pressure.


Compound of EC ₅₀ Example No.	(μ M)	maximum effect (%)
1c	9	+52 at 30 μ M
3c	4	+63 at 10 μ M
5c	>10	+14 at 30 μ M
6c	0.5	+25 at 10 μ M
7c	2.5	+29 at 10 μ M
8g	2	+64 at 10 μ M
9d	5	+50 at 30 μ M
12	5	+22 at 10 μ M
13d	10	+48 at 30 μ M
14c	1.5	+25 at 10 μ M
15c	3	+37 at 10 μ M
16c	10	+57 at 30 μ M
18e	10	+35 at 30 μ M
19e	6	+39 at 30 μ M

EXAMPLE 4.

The preparation of PLB deactivators is described below by the following non-limiting examples.

Example 1. Preparation of 3-Benzyl-5,7-bis[(1*H*-tetrazol-5-yl)-methoxy]-4-methyl-2*H*-1-benzopyran-2-one


a) 3-Benzyl-5,7-dihydroxy-4-methyl-2*H*-1-benzopyran-2-one



A solution of phloroglucinol dihydrate (20 g) and ethyl 2-benzylaceto-acetate (27.5 ml) in ethanol (320 ml) was treated with dry HCl at 0°C for five hours and the solution was kept at that temperature overnight. The yellow solution was concentrated and triturated with water, the solids filtered, washed with water and dried. The resulting hydrate was thrice evaporated to dryness from toluene, triturated with pethroleum ether (bp. 40-60°C) and filtered. Yield 33,4 g (96 %). Melting point 258-260 °C.

¹H-NMR (DMSO-d₆, 400MHz): 2.525 (s, 3H, CH₃), 3.887 (s, 2 H, CH₂Ph), 6.171 (d, 1H, J = 2,4 Hz), 6.274 (d, 1H, J = 2,4 Hz), 7.167-7.279 (m, 5H, Ph), 10.2 (s, 1H, OH), 10.47 (s, 1H, OH).


b) 3-Benzyl-5,7-bis(cyanomethoxy)-4-methyl-2*H*-1-benzopyran-2-one



Chloroacetonitrile (6.86 g), potassium carbonate (23.9 g) and 12.2 g of the product from example 1a were stirred in 120 ml of DMF at 100°C under nitrogen for two hours. The reaction mixture was cooled and poured into ice water. The solids were filtered and washed with water. Yield 13.8 g (88 %). Melting point 147-154°C.

¹H-NMR (DMSO-d₆, 400MHz): 2.525 (s, 3H, CH₃), 3.969 (s, 2H, CH₂Ph), 5.307 (s, 2H, OCH₂CN), 5.314 (s, 2H, OCH₂CN), 6.814 (d, 1 H, J = 2.5 Hz), 6.940 (d, 1H, J = 2.5 Hz), 7.18-7.292 (m, 5H, Ph).

c) 3-Benzyl-5,7-bis[(1H -tetrazol-5-yl)methoxy]-4-methyl-2H -1-benzopyran-2-one




The product from example 1b (1 g), sodium azide (0.42 g) and ammonium chloride (0.34 g) were stirred in DMF (5 ml) under nitrogen at 100 °C for 5 hours. The reaction mixture was allowed to cool down and then poured into ice water. The pH of the solution was adjusted to 10-11 and then the solution either extracted once with ethyl acetate or filtered through CELITE. The solution was acidified to pH 2 with hydrochloric acid, kept at 5°C and filtered. Yield 0.96 g (81 %). Melting point 229-233°C.

¹H-NMR (DMSO-d₆, 400MHz): 2.468 (s, 3H, CH₃), 3.937 (s, 2H, CH₂Ph), 5.596 (s, 2H, OCH₂Tet), 5.602 (s, 2H, OCH₂Tet), 6.832 (d, 1H, J = 2.4 Hz), 6.851 (d, 1H, J = 2.4 Hz), 7.171-7.283 (m, 5H, Ph).

Example 2. Preparation of 7,8,9,10-Tetrahydro-1,3-bis[(1*H*-tetrazol-5-yl)methoxy]-7-phenyl-6H-dibenzo[b,d]pyran-6-one


a) 7,8,9,10-Tetrahydro-1,3-dihydroxy-7-phenyl-6H-dibenzo[b,d]pyran-6-one



A solution of phloroglucinol (0.7 g) and 2-ethoxycarbonyl-3-phenylcyclohexanone (1.5 g) in ethanol was treated with dry HCl as described in example 1a. The product was first recrystallized from ethanol-water (1:1) and then triturated with ether. Yield 0.61 g.

¹H-NMR (DMSO-d₆, 400MHz): 1.38-1.52 (m, 1H), 1.57-1.66 (m, 1H), 1.69-1.78 (m, 1H), 1.86-1.96 (m, 1H), 2.9-3.02 (m, 1H), 3.3-3.4 (m, 1H), 4.050 (b, 1H), 6.157 (d, 1H, J = 2.4 Hz), 6.297 (d, 1H, J = 2.4 Hz), 7.076-7.265 (m, 5H), 10.153 (s, 1H), 10.456 (s, 1H).


b) 7,8,9,10-Tetrahydro-1,3-bis(cyanomethoxy)-7-phenyl-6H-dibenzo[b,d]pyran-6-one



The product from example 2a (0.5 g) was treated with chloroaceto-nitrile (0.25 g) and potassium carbonate (1.12g) in DMF (5 ml) as described in example 1b. Yield 0.6 g.

¹H-NMR (DMSO-d₆, 400MHz): 1.38-1.58 (m, 1H), 1.6-1.7 (m, 1H), 1.7-1.76 (m, 1H), 1.89-1.99 (m, 1H), 2.9-3.03 (m, 1H), 3.2-3.28 (m 1H), 4.111(b, 1H), 5.314 (s, 2H), 5.349 (s, 2H), 6.840 (d, 1H, J = 2.5 Hz), 6.925 (d, 1H, J = 2.5 Hz), 7.108-7.274 (m, 5H).

c) 7,8,9,10-Tetrahydro-1,3-bis[(1*H* -tetrazol-5-yl)methoxy]- 7-phenyl- 6H-dibenzo[b,d]pyran-6-one




The product from example 2b (0.6 g) was treated with sodium azide (0.2 g) and ammonium chloride (0.17 g) in DMF (5 ml) as in example 1c. The product was recrystallized from a mixture of DMF, ethanol and water (approximately 1:2:3). Yield 0.41 g. Melting point: 153-154°C.

¹H-NMR (DMSO-d₆, 400MHz): 1.38-1.5 (m, 1H), 1.5-1.6 (m, 1H), 1.69-1.76 (m, 1H), 1.87-1.96 (m, 1H), 2.9-3.05 (m, 1H), 3.2-3.3 (m, 1H), 4.094 (b, 1H), 5.602 (s,

2H), 5.643 (s, 2H), 6.832 (d, 1H, $J = 2.3$ Hz), 6.851 (d, 1H, $J = 2.3$ Hz), 7.089-7.212 (m, 5H).

Example 3. Preparation of 3-Benzyl-5,7-bis[(2,5-dihydro-5-oxo-4H-1,2,4-oxadiazol-3-yl)-methoxy]-4-methyl-2H-1-benzopyran-2-one


a) 3-Benzyl-5,7-bis[(hydroxyamidino)methoxy]-4-methyl-2H-1-benzopyran-2-one



Triethylamine (1.94 ml) was added to a suspension of hydroxylamine hydrochloride (0.97 g) in DMSO (2 ml) and the resulting mixture stirred at room temperature for thirty minutes. The crystals were filtered and washed with THF. The filtrate was concentrated and the product from example 1b (0.5 g) added. This solution was kept at 75 °C overnight. The reaction mixture was treated with ice water, the pH adjusted to 11 and the solids filtered, washed with water, and dried. Yield 0.5 g. Melting point: 155-160°C.

$^1\text{H-NMR}$ (DMSO- d_6 , 400MHz): 2.56 (s, 3H, CH_3), 3.938 (s, 2H), 4.466 (s, 2H), 4.486 (s, 2H), 5.565 (s, H, NH_2), 5.709 (s, 2H, NH_2), 6.658 (d, 1H, $J = 2.4$ Hz), 6.692 (d, 1H, $J = 2.4$ Hz), 7.168-7.284 (m, 5H, Ph), 9.346 (s, 1H, OH), 9.362 (s, 1H, OH).


b) 3-Benzyl-5,7-bis[(ethoxycarbonyloxyamidino)methoxy]-4-methyl-2H-1-benzopyran-2-one



Ethyl chloroformate (0.45 ml) was added to a solution of the product from example 3a (1 g) and pyridin (0.38 ml) in DMF (5 ml) at 0°C. The reaction mixture was kept at that temperature for an additional 30 minutes and then ice water added. The solids were filtered and washed with water. Yield 1.63 g. Melting point 87-92°C.

¹H-NMR (DMSO-d₆, 400MHz): 1.215-1.256 (m, 6H), 2.553 (s, 3H), 3.947 (s, 2H), 4.140-4.198 (m, 4H), 4.566 (s, 2H), 4.599 (s, 2H), 6.688 (d, 1H, J = 2.4 Hz), 6.718 (d, 1H, J = 2.4 Hz), 6.792 (b, 2H, NH₂), 6.818 (b, 2H, NH₂), 7.171-7.285 (m, 5H).

c) 3-Benzyl-5,7-bis[(2,5-dihydro-5-oxo-4H -1,2,4-oxadiazol-3-yl)- methoxy]-4-methyl-2H -1-benzopyran-2-one




The product from the previous example (1.5 g) and DBU (0.8 ml) in DMF (5 ml) was stirred at room temperature overnight. The reaction mixture was treated with ice water and acidified. The solids were filtered and washed with water. The resulting solid mass was taken in 0.1 N sodium hydroxide solution, treated with activated carbon and finally acidified. Yield 0.64 g. Melting point: 130-136°C.

$^1\text{H-NMR}$ (DMSO- d_6 , 400MHz): 2.524 (s, 3H), 3.954 (s, 2H), 5.187 (s, 2H), 5.215 (s, 2H), 6.748 (d, 1H, $J = 2.4$ Hz), 6.834 (d, 1H, $J = 2.4$ Hz), 7.158-7.289 (m, 5H), 12.8 (b, 2H).

Example 4. Preparation of 7,8,9,10-Tetrahydro-bis[(1*H*-tetrazol-5-yl)methoxy] - 1,3-dihydroxy-6*H*-dibenzo[b,d]pyran-6-one


a) 7,8,9,10-Tetrahydro-1,3-dihydroxy-6*H*-dibenzo[b,d]pyran-6-one



Phloroglucinol (1 g) and ethyl 2-oxocyclohexane carboxylate (1.32 g) were stirred in 75 % sulfuric acid (10 ml) overnight, the mixture poured into ice water and filtered. Yield: 1.55 g.

$^1\text{H-NMR}$ (DMSO- d_6 , 400MHz): 1.65 (b, 4H), 2.345 (b, 2H), 3.037 (b, 2H), 6.138 (d, 1H, $J = 2.4$ Hz), 6.245 (d, 1H, $J = 2.4$ Hz), 10.069 (b, 1H, OH), 10.322 (s, 1H, OH).


b) 7, 8, 9, 10-Tetrahydro-bis(cyanomethoxy)-1,3-dihydroxy-6*H*-dibenzo[b,d]pyran-6-one



The product from the previous example (0.5 g), chloroacetonitrile (0.34 g) and potassium carbonate (1.5 g) in DMF (5 ml) were reacted as in example 1b. Yield: 0.44 g.

$^1\text{H-NMR}$ (DMSO- d_6 , 400MHz): 1.68 (b, 4H), 2.41 (b, 2H), 3.00 (b, 2H), 5.297 (s, 2H), 5.309 (s, 2H), 6.797 (d, 1H, $J = 2.4$ Hz), 6.899 (d, 1H, $J = 2.4$ Hz).

c) 7,8,9,10-Tetrahydro-bis[(1*H*-tetrazol-5-yl)methoxy] -1,3-dihydroxy-6*H*-dibenzo[b,d]pyran-6-one




The product from the previous example (0.4 g) was treated with sodium azide (0.18 g) and ammonium chloride (0.14 g) in DMF (2.5 ml) as in example 1c. The product was recrystallized from ethanol-DMF (1:1). Yield 0.17 g. Melting point 283-286°C.

$^1\text{H-NMR}$ (DMSO- d_6 , 400MHz): 1.626 (b, 4H), 2.393 (b, 2H), 2.971 (b, 2H), 5.583 (s, 2H), 5.599 (s, 2H), 6.811 (s, 2H).

Example 5. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-phenyl-2*H*-1-benzopyran-2-one


a) 5,7-Dihydroxy-4-phenyl-2*H*-1-benzopyran-2-one



A solution of phloroglucinol (2.00 g) and ethyl benzoylacetate (3.05 g) in ethanol (30 ml) was treated with dry HCl as described in example 1a. The product was recrystallized from ethanol-water (1:1). Yield 3.0 g (75 %).

$^1\text{H-NMR}$ (DMSO- d_6 , 300 MHz): 5.739 (s, 1H, CH=C), 6.155 (d, 1H, $J = 2.3$ Hz), 6.263 (d, 1H, $J = 2.3$ Hz), 7.305-7.381 (m, 5H, Ph), 10.084 (s, 1H, OH), 10.368 (s, 1H, OH).


b) 5,7-Bis(cyanomethoxy)-4-phenyl-2*H*-1-benzopyran-2-one



The product from previous example (1.00 g) was treated with chloroaceto-nitrile (0.62 g) and potassium carbonate (2.72 g) in DMF (5 ml) as described in example 1b. The reaction mixture was poured into ice water and the mixture extracted with ethyl acetate. Ethyl acetate was washed with 1 M NaOH, dried with sodium sulfate and evaporated. The product was recrystallized from isopropanol. Yield 0.41 g (31 %).

$^1\text{H-NMR}$ (DMSO- d_6 , 300 MHz): 4.845 (s, 2H, OCH_2CN), 5.344 (s, 2H, OCH_2CN), 6.086 (s, 1H, CH=C), 6.770 (d, 1H, $J = 2.4$ Hz), 7.040 (d, 1H, $J = 2.4$ Hz), 7.320-7.443 (m, 5H, Ph).

c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-phenyl-2*H*-1-benzopyran-2-one




The product from previous example (0.40g) was treated with sodium azide (0.16 g) and ammonium chloride (0.14 g) in DMF (2 ml) at 100 °C for 2 hours. The product was isolated as described in example 1c. Yield: 0.40 g (79 %). Melting point 222-224 °C.

¹H-NMR (DMSO-d₆, 400 MHz): 5.148 (s, 2H, OCH₂Tet), 5.649 (s, 2H, OCH₂Tet), 5.968 (s, 1H, CH=C), 6.811 (d, 1H, J = 2.3 Hz), 6.962 (d, 1H, J = 2.3 Hz), 6.994-7.185 (m, 5H, Ph).

Example 6. Preparation of 7,8,9,10-Tetrahydro-1,3-bis[(1*H*-tetrazol-5-yl)methoxy]-8-phenyl-6*H*-dibenzo[*b,d*]pyran-6-one


a) 7,8,9,10-Tetrahydro-1,3-dihydroxy-8-phenyl-6*H*-dibenzo[*b,d*]pyran-6-one



A solution of phloroglucinol (1.56 g) and ethyl 2-oxo-5-phenylcyclo-hexane-carboxylate (2.52 g) in ethanol (25 ml) was treated with dry HCl as described in example 1a. The precipitate was filtered and washed with water and EtOH. Yield 1.0 g (32 %).

¹H-NMR (DMSO-d₆, 400 MHz): 1.72-1.82 (m, 1H), 2.01 (b, 1H), 2.317-2.387 (m, 1H), 2.707-2.763 (m, 1H), 2.830 (b, 1H), 3.041 (b, 1H), 3.35 and 3.40 (b, 1H), 6.174 (d, 1H, J = 2.3 Hz), 6.277 (d, 1H, J = 2.3 Hz), 7.200-7.350 (m, 5H, Ph), 10.131 (s, 1H, OH), 10.401 (s, 1H, OH).


b) 7,8,9,10-Tetrahydro-1,3-bis(cyanomethoxy)-8-phenyl-6*H*-dibenzo[*b,d*]pyran-6-one



The product from previous example (1.0 g) was treated with chloro-acetonitrile (0.57 g) and potassium carbonate (1.0 g) in DMF (5 ml) as described in example 1b. DMF was evaporated and residue dissolved in EtOAc. Ethyl acetate was washed with 1 M NaOH, dried with sodium sulfate and evaporated. The product was recrystallized from acetone-isopropanol (1:3). Yield 0.50 g (40 %).

¹H-NMR (DMSO-d₆, 300 MHz): 1.75-1.88 (m, 1H), 2.05 (b, 1H), 2.38-2.48 (m, 1H), 2.77-2.85 (m, 1H), 2.90 (b, 1H), 3.07 (b, 1H), 3.22 and 3.28 (b, 1H), 5.316 (s, 2H, OCH₂CN), 5.331 (s, 2H, OCH₂CN), 6.829 (d, 1H, J = 2.5 Hz), 6.939 (d, 1H, J = 2.5 Hz), 7.210-7.380 (m, 5H, Ph).

c) 7,8,9,10-Tetrahydro-1,3-bis[(1*H*-tetrazol-5-yl)methoxy]-8-phenyl-6*H*-dibenzo[b,d]pyran-6-one




The product from previous example (0.30 g) was treated with sodium azide (0.10 g) and ammonium chloride (0.09 g) in DMF (2 ml) at 100 °C for 3.5 hours. The product was isolated in the same manner as in example 1c. Yield 0.30 g (82 %). Melting point 235-245 °C.

$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz): 1.70-1.80 (m, 1H), 1.96 (b, 1H), 2.38-2.446 (m, 1H), 2.836 (m, 2H), 3.052 (b, 1H), 3.252 and 3.301 (b, 1H), 5.604 (s, 2H, OCH_2CN), 5.632 (s, 2H, OCH_2CN), 6.827 (d, 1H, $J = 2.5$ Hz), 6.858 (d, 1H, $J = 2.5$ Hz), 7.209-7.351 (m, 5H, Ph).

Example 7. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-methyl-3-(2-phenylethyl)-2*H*-1-benzopyran-2-one


a) 5,7-Dihydroxy-4-methyl-3-(2-phenylethyl)-2*H*-1-benzopyran-2-one



A solution of phloroglucinol (0.87 g) and ethyl 2-(2-phenylethyl)-acetoacetate (1.62 g) in ethanol (30 ml) was treated with dry HCl as described in example 1a. Yield: 1.77 g (87 %). Melting point 248-252 °C.

$^1\text{H-NMR}$ (DMSO-d_6 , 300 MHz): 2.413 (s, 3H, CH_3), 2.652-2.782 (m, 4H, CH_2CH_2), 6.151 (d, 1H, $J = 2.4$ Hz), 6.256 (d, 1H, $J = 2.4$ Hz), 7.183-7.304 (m, 5H, Ph), 10.137 (s, 1H, OH), 10.369 (s, 1H, OH).


b) 5,7-Bis(cyanomethoxy)-4-methyl-3-(2-phenylethyl)-2*H*-1-benzopyran-2-one



The product from previous example (0.90 g) was treated with chloroacetonitrile (0.48 g) and potassium carbonate (2.1 g) in DMF (5 ml) at 100 °C for 0.5 hours. The product was isolated as described in example 1b. Yield 1.00 g (88 %). Melting point 179-183 °C.

$^1\text{H-NMR}$ (DMSO- d_6 , 300 MHz): 2,384 (s, 3H, CH_3), 2.699-2,754 (m, 2H, CH_2CH_2), 2.805-2.841 (m, 2H, CH_2CH_2), 5,302 (s, 4H, OCH_2CN), 6,790 (d, 1H, $J = 2.5$ Hz), 6.909 (d, 1H, $J = 2.5$ Hz), 7.190-7.307 (m, 5H, Ph).

c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-methyl-3-(2-phenylethyl)-2*H*-1benzopyran-2-one




The product from previous example (0.40g) was treated with sodium azide (0.15 g) and ammonium chloride (0.12 g) in DMF (2 ml) at 100 °C for 2.5 hours. The product was isolated as described in example 1c. Yield 0.385 g (78 %). Melting point 248-250 °C.

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): 2.368 (s, 3H, CH_3), 2.668-2.707 (m, 2H, CH_2CH_2), 2.783-2.822 (m, 2H, CH_2CH_2), 5.593 (s, 2H, OCH_2Tet), 5.604 (s, 2H, OCH_2Tet), 6.819 (d, 1H, $J = 2.3$ Hz), 6.834 (d, 1H, $J = 2.3$ Hz), 7.161-7.291 (m, 5H, Ph).

Example 8. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-1,3-dibenzyl-4-methyl-2(1*H*)-quinolinone


a) 2-Benzyl-3-oxobutanoic acid 3,5-dimethoxyanilid



3,5-Dimethoxyaniline (5 g) was added in portions to a preheated (160 °C) ethyl 2-benzyl acetoacetate (15 ml) under nitrogen and kept at that temperature for 60 minutes. The cooled solution was diluted with heptane-ethyl ether and filtered. Yield 5.2 g (49 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.183 (s, 3H), 3.069 (d, 2H, J = 7.2 Hz), 3.923 (t, 1H, J = 7.2 Hz), 6.616 (dd, 1H, J = 2.3 Hz), 6.765 (d, 2H, J = 2.3 Hz), 7.13-7.3 (m, 5H), 10.123 (s, 1H).


b) 3-Benzyl-5,7-dimethoxy-4-methyl-2(1H)-quinolinone



The product from the previous example (1.2 g) was added to a preheated (85 °C) methanesulfonic acid (3.5 ml) and kept at that temperature for 15 minutes. The solution was allowed to cool and then treated with ice water. The product was filtered, washed with sodium bicarbonate and water. Yield 1.08 g (95 %).

¹H-NMR (300 MHz): 2.486 (s, 3H), 3.785 (s, 3H), 3.808 (s, 3H), 3.985 (s, 2H), 6.315 (d, 1H, J = 2.4 Hz), 6.472 (d, 1H, J = 2.4 Hz), 7.1-7.3 (m, 5 H), 11.52 (s, 1H).


c) 3-Benzyl-5,7-dihydroxy-4-methyl-2(1H)-quinolinone



The product from the previous example (1 g) was refluxed under nitrogen in pyridine hydrochloride (5 g) for twenty minutes. The reaction mixture was treated with water and the product filtered. Yield 0.9 g (100 %). Melting point: 307 - 312 °C.

¹H-NMR (300 MHz): 2.503 (s, 3H), 3.942 (s, 2H), 6.102 (d, 1H, J = 2.3 Hz), 6.187 (d, 1H, J = 2.3 Hz), 7.1-7.25 (m, 5H), 9.725 (s, 1H), 9.984 (s, 1H), 11.285 (s, 1H).


d) 1,3-Dibenzyl-5,7-dimethoxy-4-methyl-2(1H)-quinolinone



The product from the example 8b (1 g), potassium t-butoxide (0.62 g) and benzyl bromide (0.68 ml) were stirred in DMSO (10 ml) at 60 °C for 4 hours. The reaction mixture was treated with water, extracted with toluene and evaporated. The product was triturated with ethyl ether and filtered. Yield 0.5 g (39 %).

¹H-NMR (400 MHz): 2.537 (s, 3H), 3.708 (s, 3H), 3.826 (s, 3H), 4.124 (s, 2H), 5.56 (b, 2H), 6.413-6.434 (m, 2H), 7.154- 7.332 (m, 10H).


e) 1,3-Dibenzyl-5,7-dihydroxy-4-methyl-2(1H)-quinolinone.



The product from the previous example (2 g) was treated with pyridine hydrochloride (10 g) as described in example 8c. The product was extracted with ethyl acetate and evaporated. Yield 1,4 g (75 %).

¹H-NMR (400 MHz): 2.570 (s, 3H), 4.076 (s, 2H), 5.450 (b, 2H), 6.135 (d, 1H, J = 2.2 Hz), 6.199 (d, 1H, J = 2.2 Hz), 7.128 - 7.333 (m, 10 H), 9.83 (b, 1H), 10.166 (s, 1H).


f) 5,7-Bis(cyanomethoxy)-1,3-dibenzyl-4-methyl-2(1H)-quinolinone.



The product from the previous example (1.4 g) was treated with chloroacetonitrile (0.76 g) and K_2CO_3 (2.5 g) in DMF (20 ml) as described in example 1b. Yield 1.5 g (89 %).

1H -NMR (400 MHz): 2.555 (s, 3H), 4.146 (s, 2H), 5.214 (s, 2H), 5.275 (s, 2H), 5.578 (s, 2H), 6.735 (s, 2H), 7.13-7.33 (m, 10H).

g) 5,7-Bis[(1H-tetrazol-5-yl)methoxy]-1,3-dibenzyl-4-methyl-2(1H)-quinolinone.




The product from the previous example (1.3 g) was treated with sodium azide (0.41 g) and ammonium chloride (0.34 g) as described in example 1c. Yield: 0.69 g (45 %).

1H -NMR (400 MHz): 2.471 (s, 3H), 4.113 (s, 2H), 5.477 (s, 2H), 5.55 (b, 2H), 5.574 (s, 2H), 6.670 (d, 1H, $J = 2.1$ Hz), 6.775 (d, 1H, $J = 2.1$ Hz), 7.13-7.32 (m, 10 H).

Example 9. Preparation of 5,7-Bis[(1H-tetrazol-5-yl)methoxy]-3-benzyl-1,4-dimethyl-2(1H)-quinolinone.


a) 3-Benzyl-5,7-dimethoxy-1,4-dimethyl-2(1H)-quinolinone.



The product from example 8b (0.5 g), t-BuOK (0.2 g) and methyl iodide (0.4 ml) were stirred in DMSO (5 ml) at 35 °C for two days. The reaction mixture was treated with water and extracted with toluene. The product was purified by column chromatography using toluene-ethyl acetate-acetic acid 8 : 2: 1 as the eluent. Yield 0.24 g (46 %).

¹H-NMR (300 MHz): 2.51 (s, 3H), 3.632 (s, 2H), 3.846 (s, 3H), 3.896 (s, 3H), 4.047 (s, 2H), 6.468 (d, 1H, J = 2.3 Hz), 6.558 (d, 1H, J = 2.3 Hz), 7.1-7.26 (m, 5H).


b) 3-Benzyl-5,7-dihydroxy-1,4-dimethyl-2(1H)-quinolinone.



The product from the previous example (0.2 g) was treated with pyridine hydrochloride (2 g) as described in example 8c and the product extracted with ethyl acetate. Yield 0.16 g (89 %).

¹H-NMR (400 MHz): 2.567 (s, 3H), 3.515 (s, 3H), 4.005 (s, 2H), 6.244 (d, 1H, J = 2.3 Hz), 6.268 (d, 1H, J = 2.3 Hz), 7.08-7.25 (m, 5H), 9.879 (s, 1H), 10.113 (s, 1H).


c) 5,7-Bis(cyanomethoxy)-3-benzyl-1,4-dimethyl-2(1H)-quinolinone.



The product from the previous example (0.15 g), chloroacetonitrile 0.08 g) and K_2CO_3 (0.28 g) were reacted in DMF (2 ml) as described in example 1b. Yield 0.16 g (84 %).

1H -NMR (400 MHz): 2.524 (s, 3H), 3.658 (s, 3H), 4.079 (s, 2H), 5.292 (s, 2H), 5.379 (s, 2H), 6.766 (d, 1H, $J = 2.3$ Hz), 6.855 (d, 1H, $J = 2.3$ Hz), 7.13-7.24 (m 5H).


d) 5,7-Bis[(1H-tetrazol-5-yl)methoxy] -3-benzyl-1,4-dimethyl-2(1H)-quinolinone.



The product from the previous example (0.15 g) was treated with NaN_3 (57 mg) and NH_4Cl (47 mg) in DMF (2 ml) as described in example 1c. Yield 0.115 g. Melting point: 250-253°C.

1H -NMR (400 MHz): 2.451 (s, 3H), 3.649 (s, 3H), 4.042 (s, 2H), 6.792 (d, 1H, $J = 2.2$ Hz), 6.833 (d, 1H, $J =$ Hz), 7.1-7.25 (m, 5H).

Example 10. Preparation of 3-Benzyl-5,7-bis[(2-methyl-1H -tetrazol-5-yl)methoxy]-4-methyl-2H -1-benzopyran-2-one and the three isomers.




0.07 ml of methyl iodide was added to a solution of 0.2 g of the product from example 1c and 0.31 g of K_2CO_3 in 2 ml of DMF and the mixture stirred at room temperature for 4 hours. The reaction mixture was poured into ice water and filtered. Yield 0.2 g as a mixture of four regioisomers, melting point 71-76°C.

1H -NMR (DMSO- d_6 , 400MHz): 2.47 (s, CH_3), 2.48 (s, CH_3), 3.93 (s, CH_2Ph), 4.11 (s, NCH_3), 4.12 (s, NCH_3), 4.15 (s, NCH_3), 4.38 (s, NCH_3), 4.40 (s, NCH_3), 5.51 (s, OCH_2), 5.52 (s, OCH_2), 5.62 (s, OCH_2), 5.67 (s, OCH_2), 6.84-6.91 (m, 2H), 7.16-7.28 (m, 5H, Ph).

Example 11. Preparation of 3-Benzyl-5,7-bis[1-(1 *H*-tetrazol-5-yl)ethoxy]4-methyl-2 *H*-1-benzopyran-2-one, mixture of stereoisomers


a) 3-Benzyl-5,7-bis-[(1-cyano)ethoxy]-4-methyl-2 *H*-1-benzopyran-2-one



The product from example 1a (1 g), 2-chloropropionitrile (0.7 g) and potassium carbonate (2 g) were heated in DMF (15 ml) under nitrogen at 110°C for sixty minutes. The mixture was treated with water, filtered and washed with 1 N NaOH and water. Yield 1.2 g.

1H -NMR (DMSO- d_6 , 300MHz): 1.74-1.78 (t + t, 6 H, $CH-CH_3$), 2.53 (s, 3 H), 3.97 (s, 2H), 5.58-5.66 (m, 2H, $CH-CH_3$), 6.87 (m, 1H), 6.99 (d, 1H), 7.18-7.31 (m, 5H).


b) 3-Benzyl-5,7-bis[1-(1 *H* -tetrazol-5-yl)ethoxy]4-methyl-2 *H* -1-benzopyran-2-one, mixture of stereoisomers.



The product from the previous example (0.5 g), sodium azide (0.18 g) and ammonium chloride (0.15 g) were heated in DMF (7 ml) at 100 °C for 90 minutes. The product was treated with water, extracted with ethyl acetate and evaporated. Yield 0.57 g. Melting point 91-104°C.

¹H-NMR (DMSO-*d*₆, 300MHz): 1.69-1.77 (m, 6 H, CH-CH₃), 2.54 (s, 3H), 3.94 (s, 2H), 6.10-6.17 ((m, 2H, CH-CH₃), 6.65 (dd, 1H), 6.74 (dd, 1H), 7.13-7.30 (m, 5H).

Example 12. Preparation of 5,7-Bis(carboxymethoxy)-1,3-dibenzyl-4-methyl-2(1H)-quinolinone




The product from example 8f (0.2 g) was refluxed in a solution of concentrated hydrochloric acid (3 ml) and acetic acid (2 ml) for one hour. The product was filtered at 25 °C. Yield 0.14 g.

¹H-NMR (300 Mhz, DMSO-d₆): 2.63 (s, CH₃), 4.14 (s, 2H, CH₂Ph), 4.66 (s, 2 H, OCH₂COOH), 4.79 (s, 2H, OCH₂COOH), 5.53 (s, 2H, NCH₂Ph), 6.41 (d, 1H, J = 2.2 Hz), 6.45 (d, 1H, J = 2.2 Hz), 7.13-7.34 (m, 10 H, Ph).

Example 13. Preparation of 3-Benzyl-5,7-bis[(1H-tetrazol-5-yl)methoxy]-1-(4-fluorobenzyl)-4-methyl-2(1H)-quinolinone


a) 1-Benzyl-5,7-dimethoxy-3-(4-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from example 8b (2 g), potassium -tert-butoxide (0.87 g) and 4-fluorobenzylchloride (1.12 g) were heated in DMSO (20 ml) at 60 °C for three hours as in example 8d. Yield 1.28 g.

¹H-NMR (400 Mhz, DMSO-d₆): 2.53 (s, 3H), 3.73 (s, 3H), 3.83 (s, 3H), 5.55 (s, 2H), 6.43 (s, 2H), 7.12-7.2 (m, 5 H), 7.26-7.28 (m, 4H).


b) 3-Benzyl-5,7-dihydroxy-1-(4-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from previous example (1.25 g) were heated in pyridine hydrochloride (12.5 g) at about 225 °C for 9 minutes. Yield 1 g.

¹H-NMR (300 Mhz, DMSO-d₆): 2.56 (s, 3H), 4.07 (s, 2H), 5.4 (b, 2H), 6.13 (d, 1H, J = 2.1 Hz), 6.20 (d, 1H, J = 2.1 Hz), 7.12-7.28 (m, 9H), 9.88 (s, 1H), 10.22 (s, 1H).


c) 3-Benzyl-5,7-Bis(cyanomethoxy)-1-(4-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from the previous example (1 g), ClCH₂CN (0.43 g) and K₂CO₃ (1.42 g) were heated in DMF (8 ml) at 120 °C for one hour. Yield 0.94 g.

¹H-NMR (300 Mhz, DMSO-d₆): 2.55 (s, 3H), 4.14 (s, 2H), 5.25 (s, 2H), 5.28 (s, 2H), 5.57 (s, 2H), 6.74 (s, 2H, ArH), 7.1 -7.3 (m, 9H).

d) 3-Benzyl-5,7-bis[(1H-tetrazol-5-yl)methoxy]-1-(4-fluorobenzyl)-4-methyl-2(1H)-quinolinone




The product from the previous example (0.5 g), sodium azide (0.14 g) and ammonium chloride (0.12 g) were heated in DMF (5 ml) at 120 °C for 90 min. The product was triturated with acetonitrile. Yield 0.28 g. Melting point: 126-132 °C.

¹H-NMR (300 Mhz, DMSO-d₆): 2.48 (s, 3H), 4.11 (s, 2H), 5.51 (s, 2H), 5.55 (s, 2H), 5.58 (s, 2H), 6.67 (d, 1H, J = 2.1 Hz), 6.78 (d, 1H, J = 2.1 Hz).

Example 14. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-(4-chlorobenzyl)-4-methyl-2*H*-1-benzopyran-2-one


a) 3-(4-Chlorobenzyl)-5,7-dihydroxy-4-methyl-2*H*-1-benzopyran-2-one



A solution of phloroglucinol (1.57 g) and ethyl 2-(4-chlorobenzyl)-acetoacetate (3.18 g) in ethanol (25 ml) was treated with dry HCl at 0 °C for 1.5 hours and the solution was kept at that temperature overnight. Solvent was evaporated and the precipitate triturated with water. Yield 3.87 g (98 %). Melting point 270-278 °C.

¹H-NMR (DMSO-d₆, 300 MHz): 2.52 (s, 3H, CH₃), 3.87 (s, 2H, CH₂), 6.17 (d, 1H, J = 2.4 Hz), 6.28 (d, 1H, J = 2.4 Hz), 7.18-7.34 (m, 4H, Ph), 10.21 (s, 1H, OH), 10.48 (s, 1H, OH).


b) 5,7-Bis(cyanomethoxy)-3-(4-chlorobenzyl)-4-methyl-2*H*-1-benzopyran-2-one



The product from the previous example (1.00 g), chloroacetonitrile (0.50 g) and potassium carbonate (2.18 g) were heated in DMF (5 ml) at 100 °C for 30 minutes. The product was isolated as described in example 1b. Yield 0.90 g (72 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.52 (s, 3H, CH₃), 3.95 (s, 2H, CH₂), 5.308 (s, 2H, OCH₂CN), 5.312 (s, 2H, OCH₂CN), 6.81 (d, 1H, J = 2.5 Hz), 6.94 (d, 1H, J = 2.5 Hz), 7.22-7.33 (m, 4H, Ph).


c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-(4-chlorobenzyl)-4-methyl-2*H*-1-benzopyran-2-one



The product from the previous example (0.40 g), sodium azide (0.14 g) and ammonium chloride (0.11 g) were heated in DMF (2 ml) at 100 °C for 2 hours. The product was isolated as in example 1c. Yield 0.40 g (82 %).


¹H-NMR (DMSO-d₆, 300 MHz): 2.46 (s, 3H, CH₃), 3.92 (s, 2H, CH₂), 5.602 (s, 2H, OCH₂Tet), 5.609 (s, 2H, OCH₂Tet), 6.83 (d, 1H, J = 2.5 Hz), 6.85 (d, 1H, J = 2.5 Hz), 7.20-7.33 (m, 4H, Ph).

Example 15. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-(4-nitrobenzyl)-4-methyl-2*H*-1-benzopyran-2-one

a) 5,7-Dihydroxy-4-methyl-3-(4-nitrobenzyl)-2*H*-1-benzopyran-2-one

A solution of phloroglucinol (0.48 g) and ethyl 2-(4-nitrobenzyl)aceto-acetate (1.00 g) in ethanol (150 ml) was treated with dry HCl at 0 °C for 7.5 hours and the solution was kept at that temperature overnight. Solvent was evaporated and the precipitate triturated with water. Yield 0.63 g (51 %). Melting point 280-285 °C.


¹H-NMR (DMSO-*d*₆, 300 MHz): 2.53 (s, 3H, CH₃), 4.03 (s, 2H, CH₂), 6.19 (d, 1H, *J* = 2.4 Hz), 6.29 (d, 1H, *J* = 2.4 Hz), 7.40-7.51 and 8.11-8.17 (m, 4 H, Ph), 10.25 (s, 1H, OH), 10.52 (s, 1H, OH).

b) 5,7-Bis(cyanomethoxy)-3-(4-nitrobenzyl)-4-methyl-2*H*-1-benzopyran-2-one

The product from the previous example (0.57 g), chloroacetonitrile (0.27 g) and potassium carbonate (1.20 g) were heated in DMF (2 ml) at 100 °C for 50 minutes. The product was isolated as described in example 1b. Yield 0.47 g (67 %). Melting point 178-185 °C.

¹H-NMR (DMSO-*d*₆, 400 MHz): 2.53 (s, 3H, CH₃), 4.11 (s, 2H, CH₂), 5.319 (s, 2H, OCH₂CN), 5.323 (s, 2H, OCH₂CN), 6.83 (d, 1H, *J* = 2.4 Hz), 6.96 (d, 1H, *J* = 2.4 Hz), 7.48-7.53 and 8.12-8.16 (m, 4H, Ph).

c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-(4-nitrobenzyl)-4-methyl-2*H*-1-benzopyran-2-one




The product from the previous example (0.38 g), sodium azide (0.12 g) and ammonium chloride (0.11 g) were heated in DMF (3 ml) at 100 °C for 2 hours. The product was isolated as described in example 1c. Yield 0.25 g (54 %). Melting point 240-244 °C.

¹H-NMR (DMSO-*d*₆, 400 MHz): 2.47 (s, 3H, CH₃), 4.08 (s, 2H, CH₂), 5.611 (s, 2H, OCH₂Tet), 5.623 (s, 2H, OCH₂Tet), 6.85 (d, 1H, *J* = 2.4 Hz), 6.87 (d, 1H, *J* = 2.4 Hz), 7.46-7.50 and 8.12-8.16 (m, 4H, Ph).

Example 16. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-cyclopentyl-4-methyl-2*H*-1-benzopyran-2-one


a) 3-Cyclopentyl-5,7-dihydroxy-4-methyl-2*H*-1-benzopyran-2-one



A solution of phloroglucinol (2.00 g) and ethyl 2-cyclopentylaceto-acetate (3.14 g) in ethanol (40 ml) was treated with dry HCl at 0 °C for 2.5 hours and the solution kept at that temperature overnight. Solvent was evaporated and the precipitate purified with flash chromatography eluting with toluene-EtOAc-AcOH (8:1:1). Yield 1.22 g (29 %).

$^1\text{H-NMR}$ (DMSO- d_6 , 300 MHz): 1.50-1.88 (m, 8H, $-(\text{CH}_2)_4-$), 2.57 (s, 3H, CH_3), 3.25 (m, 1H, CH), 6.11 (d, 1H, $J = 2.4$ Hz), 6.25 (d, 1H, $J = 2.4$ Hz), 10.25 (b, 2H, OH).


b) 5,7-Bis(cyanomethoxy)-3-cyclopentyl-4-methyl-2*H*-1-benzopyran-2-one



The product from the previous example (0.50 g), chloroacetonitrile (0.31 g) and potassium carbonate (0.61 g) were heated in DMF (2 ml) at 80 °C for 40 minutes. The product was isolated as described in example 1b. Yield 0.56 g (86 %).

$^1\text{H-NMR}$ (DMSO- d_6 , 300 MHz): 1.55-1.90 (m, 8H, $-(\text{CH}_2)_4-$), 2.56 (s, 3H, CH_3), 3.37 (m, 1H, CH), 5.29 (s, 2H, OCH_2CN), 5.31 (s, 2H, OCH_2CN), 6.75 (d, 1H, $J = 2.5$ Hz), 6.88 (d, 1H, $J = 2.5$ Hz).

c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-3-cyclopentyl-4-methyl-2*H*-1-benzopyran-2-one




The product from the previous example (0.30 g), sodium azide (0.13 g) and ammonium chloride (0.11 g) were heated in DMF (1 ml) at 100 °C for 1.5 hours. The

product was isolated as described in example 1c. Yield 0.30 g (80 %). Melting point 248-252 °C.

¹H-NMR (DMSO-d₆, 400 MHz): 1.53-1.89 (m, 8H, -(CH₂)₄-), 2.51 (s, 3H, CH₃), 3.34 (m, 1 H, CH), 5.59 (s, 2H, OCH₂Tet), 5.61 (s, 2H, OCH₂Tet), 6.80 (s, 2H).

Example 17. Preparation of 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-methyl-3-(1-naphtylmethyl)-2*H*-1-benzopyran-2-one


a) 5,7-dihydroxy-4-methyl-3-(1-naphtylmethyl)-2*H*-1-benzopyran-2-one



A solution of phloroglucinol (0.47 g) and ethyl 2-(1-naphtylmethyl)-acetoacetate (1.00 g) in ethanol (20 ml) was treated with dry HCl at 0 °C for 3 hours and the solution kept at that temperature overnight. Solvent was evaporated and the precipitate triturated with water and recrystallized from isopropanol-water (1:1). Yield 0,96 g (78 %). Melting point 275-280 °C.

¹H-NMR (DMSO-d₆, 400 MHz): 2.45 (s, 3H, CH₃), 4.32 (s, 2H, CH₂), 6.23 (d, 1H, J = 2.5 Hz), 6.32 (d, 1 H, J = 2.5 Hz), 6.97-8.25 (m, 7H, Naph), 10.26 (s, 1H, OH), 10.53 (s, 1H, OH).


b) 5,7-Bis(cyanomethoxy)-4-methyl-3-(1-naphtylmethyl)-2*H*-1-benzopyran-2-one



The product from the previous example (0.80 g), chloroacetonitrile (0.36 g) and potassium carbonate (0.66 g) were heated in DMF (4 ml) at 100 °C for 1 hour. The product was isolated as in example 1b. Yield 0.30 g (30 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.45 (s, 3H, CH₃), 4.40 (s, 2H, CH₂), 5.34 (s, 2H, OCH₂CN), 5.36 (s, 2H, OCH₂CN), 6.86 (d, 1H, J = 2.5 Hz), 7.010 (d, 1H, J = 2.5 Hz), 7.016-8.27 (m, 7H, Naph).

c) 5,7-Bis[(1*H*-tetrazol-5-yl)methoxy]-4-methyl-3-(1-naphtylmethyl)-2*H*-1-benzopyran-2-one




The product from the previous example (0.25 g), sodium azide (0.080 g) and ammonium chloride (0.072 g) were heated in DMF (2 ml) at 100 °C for 2.5 hours. The product was isolated as described in example 1c. Yield 0.11 g (36 %). Melting point 164-174 °C.

¹H-NMR (DMSO-d₆, 300 MHz): 2.40 (s, 3H, CH₃), 4.37 (s, 2H, CH₂), 5.63 (s, 2H, OCH₂Tet), 5.65 (s, 2H, OCH₂Tet), 6.87 (d, 1H, J = 2.5 Hz), 6.92 (d, 1H, J = 2.5 Hz), 6.98-8.26 (m, 7H, Naph).

Example 18. Preparation of 1-Benzyl-5,7-bis-[(1*H*-tetrazol-5-yl)-methoxy]-4-methyl-2(1*H*)-quinolinone


a) 5,7-Dimethoxy-4-methyl-2(1*H*)-quinolinone



tert-Butyl acetoacetate (1.58 g) was heated to 120 °C and 3,5-dimethoxyaniline (1.53 g) dissolved in xylene (4 ml) was added. The mixture was heated at 120-130 °C for 20 minutes and then cooled to room temperature. Methanesulfonic acid (2 ml) was added and the mixture was stirred at ambient temperature for 10 minutes. Water (40 ml) was added and the precipitate filtered and dried. Yield 1.31 g (60 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.50 (s, 3H, CH₃), 3.79 (s, 3H, OCH₃), 3.83 (s, 3H, OCH₃), 6.03 (s, 1H, CH=C), 6.31 (d, 1H, J = 2.3 Hz), 6.45 (d, 1H, J = 2.3 Hz), 11.4 (b, 1H, NH).


b) 1-Benzyl-5,7-dimethoxy-4-methyl-2(1H)-quinolinone



The product from the previous example (1.20 g) was suspended in DMSO (15 ml) and *t*-BuOK (0.68 g) and benzylbromide (1.03 g) were added. Reaction mixture was stirred at ambient temperature overnight. Water was added and the product extracted to EtOAc. EtOAc was dried and evaporated to dryness. The product was recrystallized from toluene. Yield 0.80 g (47 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.55 (d, 3H, J = 1.1 Hz, CH₃), 3.71 (s, 3H, OCH₃), 3.84 (s, 3H, OCH₃), 5.48 (b, 2H, NCH₂), 6.29 (d, 1H, J = 1.1 Hz, CH=C), 6.4 (s, 2H), 7.18-7.33 (m, 5H, Ph).


c) 1-Benzyl-5,7-dihydroxy-4-methyl-2(1H)-quinolinone



The product from the previous example (0.69 g) was dissolved to CH_2Cl_2 (14 ml) and the reaction mixture cooled to -20°C . BBr_3 (2.4 g) in CH_2Cl_2 (1M solution) was added and the mixture was allowed to warm to ambient temperature during the night. The precipitate was filtered, washed with CH_2Cl_2 and dissolved to EtOAc. EtOAc was washed with dilute HCl, dried and evaporated to dryness. Yield 0.34 g (54 %).

$^1\text{H-NMR}$ (DMSO-d_6 , 300 MHz): 2.56 (d, 3H, $J = 1.0$ Hz, CH_3), 5.33 (b, 2H, NCH_2), 6.11 (d, 1H, $J = 2.1$ Hz), 6.13 (d, 1H, $J = 1.0$ Hz, $\text{CH}=\text{C}$), 6.17 (d, 1H, $J = 2.1$ Hz), 7.12-7.34 (m, 5H, Ph), 9.90 (b, 1H, OH), 10.22 (s, 1H, OH).


d) 1-Benzyl-5,7-bis(cyanomethoxy)-4-methyl-2(1H)-quinolinone



The product from the previous example (0.34 g), chloroacetonitrile (0.13 g) and potassium carbonate (0.34 g) were heated in DMF (2 ml) at 100°C for 1.5 hours. Water was added and the precipitate filtered and dried. The product was recrystallized from isopropanol. Yield 0.20 g (46 %).

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): 2.57 (s, 3H, CH_3), 5.22 (s, 2H, OCH_2CN), 5.30 (s, 2H, OCH_2CN), 5.50 (b, 2H, NCH_2), 6.42 (s, 1H, $\text{CH}=\text{C}$), 6.70 (d, 1H, $J = 2.1$ Hz), 6.73 (d, 1H, $J = 2.1$ Hz), 7.21-7.32 (m, 5H, Ph).

e) 1-Benzyl-5,7-bis-[(1*H*-tetrazol-5-yl)methoxy]-4-methyl-2(1*H*)-quinolinone




The product from the previous example (0.20 g), sodium azide (0.072 g) and ammonium chloride (0.060 g) were heated in DMF (2 ml) at 100 °C for 3 hours. The product was isolated as described in example 1c. Yield 0.21 g (85 %). Melting point 246-249 °C.

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): 2.50 (s, 3H, CH_3), 5.48 (b, 4H, OCH_2Tet , NCH_2), 5.60 (s, 2H, OCH_2Tet), 6.34 (s, 1H, $\text{CH}=\text{C}$), 6.64 (d, 1H, $J = 1.9$ Hz), 6.77 (d, 1H, $J = 1.9$ Hz), 7.18-7.32 (m, 5H, Ph).

Example 19. Preparation of 1-Benzyl-5,7-bis[1*H*-tetrazol-5-yl)methoxy]-3-(2-fluorobenzyl)-4-methyl-2(1*H*)-quinolinone

a) 5,7-Dimethoxy-3-(2-fluorobenzyl)-4-methyl-2(1*H*)-quinolinone




Ethyl 2-(2-fluorobenzyl)acetoacetate (2.5 g) in xylene (1 ml) was heated to 150 °C and 3,5- dimethoxyaniline (1.46 g) in xylene (4 ml) was added in small portions

during 30 minutes. The reaction mixture was heated at 160 °C for 3 hours and then cooled to room temperature. Methanesulfonic acid (1.7 ml) was added and the mixture was stirred at ambient temperature for 30 minutes. Water was added and the precipitate filtered and dried. The product was triturated with warm ethanol. Yield 0.64 g (21 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.45 (s, 3H), 3.79 (s, 3H), 3.82 (s, 3H), 3.97 (s, 2H), 6.33 (d, 1H, J = 2.4 Hz), 6.48 (d, 1H, J = 2.4 Hz), 6.90-7.25 (m, 4H), 11.61 (s, 1H).


b) 1-Benzyl-5,7-dimethoxy-3-(2-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from the previous example (0.62 g) was treated with *t*-BuOK (0.23 g) and benzylbromide (0.36 g) in DMSO (12 ml) at 60 °C for 2.5 hours. The product was isolated as described in example 18b. Yield 0.39 g (49 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.51 (s, 3H), 3.72 (s, 3H), 3.84 (s, 3H), 4.11 (s, 2H), 5.55 (b, 2H), 6.433 (d, 1H, J = 2.1 Hz), 6.443 (d, 1H, J = 2.1 Hz), 6.97-7.33 (m, 9H).


c) 1-Benzyl-5,7-dihydroxy-3-(2-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from the previous example (0.34 g) was treated with BBr₃ (8.48 g) in CH₂Cl₂ (7 ml) as described in example 18c. Yield 0.30 g (82 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.55 (s, 3H), 4.06 (s, 2H), 5.40 (b, 2H), 6.13 (d, 1H, J = 2.1 Hz), 6.22 (d, 1H, J = 2.1 Hz), 6.97-7.33 (m, 9H), 10.3 (b, 2H).


d) 1-Benzyl-5,7-bis(cyanomethoxy)-3-(2-fluorobenzyl)-4-methyl-2(1H)-quinolinone



The product from the previous example (0.21 g), chloroacetonitrile (0.086 g) and potassium carbonate (0.37 g) were heated in DMF (2 ml) at 100 °C for 2 hours. The product was isolated as described in example 1b. Yield 0.18 g (71 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.53 (s, 3H), 4.13 (s, 2H), 5.23 (s, 2H), 5.29 (s, 2H), 5.57 (b, 2H), 6.746 (d, 1H, J = 2.3 Hz), 6.756 (d, 1H, J = 2.3 Hz), 7.00-7.32 (m, 9H).

e) 1-Benzyl-5,7-bis[1H-tetrazol-5-yl)methoxy]-3-(2-fluorobenzyl)-4-methyl-2(1H)-quinolinone




The product from the previous example (0.17 g), sodium azide (0.051 g) and ammonium chloride (0.042 g) were heated in DMF at 100 °C for 3 hours. The product was isolated as described in example 1c. Yield 0.17 g (85 %). Melting point 135-140 °C.

¹H-NMR (DMSO-d₆, 400 MHz): 2.46 (s, 3H), 4.10 (s, 2H), 5.48 (s, 2H), 5.51 (b, 2H), 5.59 (s, 2H), 6.68 (d, 1H, J = 2.2 Hz), 6.79 (d, 1H, J = 2.2 Hz), 6.99-7.32 (m, 9H).

Example 20. Preparation of 1-Benzyl-5,7-bis[1*H*-tetrazol-5-yl]-methoxy]-4-methyl-3-(2-phenylethyl)-2(1*H*)-quinolinone


a) 5,7-Dimethoxy-4-methyl-3-(2-phenylethyl)-2(1*H*)-quinolinone



Ethyl 2-(2-phenylethyl)acetoacetate (2.70 g) in xylene (5 ml) was treated with 3,5-dimethoxyaniline (1.60 g) at 150 °C as described in example 19a. Methanesulfonic acid (4.0 ml) was added at room temperature and the mixture heated at 80 °C for 1 hour. The product was isolated as described in example 19a. Yield 1.38 g (41 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.45 (s, 3H), 2.64-2.68 (m, 2H), 2.82-2.86 (m, 2H), 3.78 (s, 3H), 3.81 (s, 3H), 6.30 (d, 1H, J = 2.3 Hz), 6.45 (d, 1H, J = 2.3 Hz), 7.18-7.30 (m, 5H), 11.45 (s, 1H).


b) 1-Benzyl-5,7-dimethoxy-4-methyl-3-(2-phenylethyl)-2(1*H*)-quinolinone



The product from the previous example (0.61 g), *t*-BuOK (0.24 g) and benzylbromide (0.36 g) were heated in DMSO (12 ml) at 60 °C for 2 hours. The product was isolated as described in example 18b. Yield 0.31 g (40 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.51 (s, 3H), 2.73-2.77 (m, 2H), 2.96-3.00 (m, 2H), 3.70 (s, 3H), 3.83 (s, 3H), 5.55 (b, 2H), 6.40 (s, 2H), 7.17-7.33 (m, 10 H).


c) 1-Benzyl-5,7-dihydroxy-4-methyl-3-(2-phenylethyl)-2(1H)-quinolinone



The product from the previous example (0.31 g) was treated with BBr₃ (0.75 g) in CH₂Cl₂ (5 ml) as in example 18c. Yield 0.26 g (89 %).

¹H-NMR (DMSO-d₆, 300 MHz): 2.56 (s, 3H), 2.69-2.75 (m, 2H), 2.90-2.95 (m, 2H), 5.39 (b, 2H), 6.08 (d, 1H, J = 2.0 Hz), 6.19 (d, 1H, J = 2.0 Hz), 7.11-7.33 (m, 10H), 10.2 (b, 2H).


d) 1-Benzyl-5,7-bis(cyanomethoxy)-4-methyl-3-(2-phenylethyl)-2(1H)-quinolinone



The product from the previous example (0.22 g), chloroacetonitrile (0.091 g) and potassium carbonate (0.39 g) were heated at 100 °C for 2 hours. The product was isolated as in example 1b. Yield 0.20 g (76 %).

¹H-NMR (DMSO-d₆, 400 MHz): 2.50 (s, 3H), 2.73-2.77 (m, 2H), 2.98-3.02 (m, 2H), 5.21 (s, 2H), 5.29 (s, 2H), 5.56 (b, 2H), 6.70 (d, 1H, J = 2.1 Hz), 6.72 (d, 1H, J = 2.1 Hz), 7.18-7.33 (m, 10H).


e) 1-Benzyl-5,7-bis[1*H*-tetrazol-5-yl]methoxy-4-methyl-3-(2-phenylethyl)-2(1*H*)-quinolinone



The product from the previous example (0.19 g), sodium azide (0.057 g) and ammonium chloride (0.047 g) were heated in DMF at 100 °C for 3 hours. The product was isolated as described in example 1c. Yield 0.18 g (78 %). Melting point 215-218 °C.

¹H-NMR (DMSO-d₆, 400 MHz): 2.46 (s, 3H), 2.70-2.74 (m, 2H), 2.95-2.99 (m, 2H), 5.47 (s, 2H), 5.54 (b, 2H), 5.57 (s, 2H), 6.64 (d, 1H, J = 2.0 Hz), 6.77 (d, 1H, J = 2.0 Hz), 7.16-7.33 (m, 10H).


Example 21. Preparation of 5,7-Bis(aminocarbonylmethoxy)-1,3-dibenzyl-4-methyl-2(1*H*)-quinolinone.



The mixture of 5,7-dihydroxy-1,3-dibenzyl-4-methyl-2(1H)-quinolinone (0.5 g), potassium carbonate (0.9 g) and 2-chloroacetamide (0.25 g) in DMF (6.5 ml) were reacted at 100 °C for two hours. The reaction mixture was treated with ice water and filtered. The product was triturated with hot ethanol. Yield: 0.32 g. Melting point 252-253°C.

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6): 2.63 (s, 3H, CH₃), 4.13 (s, 2H, PhCH₂), 4.37 (s, 2H, OCH₂), 4.55 (s, 2H, OCH₂), 5.54 (s, 2H, NCH₂Ph), 6.40 (d, 1H, J = 2 Hz, ArH), 6.53 (d, 1H, J = 2 Hz, ArH), 7.13-7.33 (m, 10 H, Ph), 7.44 (d, 2H, J = 65 Hz, CONH₂), 7.47 (d, 2H, J = 68 Hz, CONH₂).


Example 22. Preparation of 5,7-Bis(ethoxycarbonylmethoxy)-1,3-dibenzyl-4-methyl-2(1H)-quinolinone.



The mixture of 5,7-dihydroxy-1,3-dibenzyl-4-methyl-2(1H)-quinolinone (1 g), ethyl 2-bromoacetate (0.63 ml) and potassium carbonate (1.49 g) in DMF (5 ml) was heated under nitrogen at 110 °C for three hours, poured into ice water and filtered. The resulting solid material was triturated with ether and filtered again. Yield: 1.03 g, melting point 113-116 °C.

¹H-NMR (400 MHz, DMSO-d₆): 1.15 (t, 3H, CH₃CH₂, J = 7.1 Hz), 1.20 (t, 3H, CH₃CH₂, J = 7.1 Hz), 2.63 (s, 3H, CH₃), 4.03 (q, 2H, CH₂CH₃, J = 7.1 Hz), 4.13 (s, 2H, CH₂Ph), 4.17 (q, 2H, CH₂CH₃, J = 7.1 Hz), 4.78 (s, 2H, OCH₂), 4.90 (s, 2H, OCH₂), 6.41 (d, 1H, J = 2.2 Hz), 6.44 (d, 1H, J = 2.2 Hz), 7.13-7.33 (m, 10 H, Ph).

Example 23. Preparation of 5,7-Bis(hydroxyaminocarbonylmethoxy)-1,3-dibenzyl-4-methyl-2(1H)-quinolinone



The product from the previous example (0.3 g), hydroxylamine hydrochloride (0.32 g) and 5 N NaOH (1.05 ml) were reacted in ethanol (8 ml) at 50 °C for six hours. The reaction mixture was treated with water and made basic (pH 10) and filtered. The filtrate was acidified to pH 2 and filtered. Yield: 0.2 g, melting point 121-127°C.

¹H-NMR (400 MHz, DMSO-d₆): the tautomeric forms of hydroxamic acid are seen in OCH₂-signals: 2.63 (s, 3H, CH₃), 4.13 (s, 2H, CH₂Ph), 4.41 (s, 2H, OCH₂), 4.54 (s, 2H, OCH₂), 4.64 (s, 2H, HON=C(OH)CH₂O), 4.65 (s, 2H, HON=C(OH)CH₂O), 4.77 (s, 2H, HON=C(OH)CH₂O), 4.78 (s, 2H, HON=C(OH)CH₂O), 5.54 (s, 2H, NCH₂Ph), 6.38-6.54 (m, 2H, ArH), 7.14-7.34 (m, 10 H, Ph), 9.05 (b, 2H, NOH), 10.84 (s, 1H, HONHCO), 10.88 (s, 1H, HONHCO).

EXAMPLE 5. Design of PLB inhibitors

The three dimensional structure determined for phospholamban can be used as a target for selecting compounds that bind to the protein. In order to have good affinity for phospholamban, the ligand should have steric and electrostatic complementarity with the target. Especially, good electrostatic and/or hydrogen bonding interactions should be formed with the sites S1 and S2, and good hydrophobic interactions should be formed with the sites S3 and S4. Any of the various computer programs and databases available for such purpose can be used to design compounds that fulfill these requirements. The structure-based approaches include *de Novo* design, computer-based selection of ligands that are complementary with the target and computer-aided optimization of lead molecules. The detection of the PLB binding compounds can proceed by using the following steps:

1. The target region of the protein is selected. The binding model of the effective phospholamban deactivator peptide, cP226 (SEQ ID NO: 10), can be used to define an area on the phospholamban surface which can function as a target for phospholamban deactivators. Especially, this determines the side chains of phospholamban which can interact with the compounds to be designed.

2. Small molecules which are complementary to the binding site can be docked to the target by using available software, such as e.g. Ludi, DOCK or LeapFrog. Computer databases of three-dimensional structures of small molecules or molecular fragments can be used in the docking. Such an approach gives molecules or fragments that have good interactions with the various parts of the target area.

3. Different small molecules or fragments that bind to the target area can be linked together or one can incorporate new side chains and/or functional groups to them, so that one gets a single, larger molecule. The resulting new compound is likely to have better affinity to the target than the smaller molecules.

4. One can also select a proper scaffold and dock that by using an interactive molecular graphics system near the binding site of the protein. One can then add new fragments and functional groups to the scaffold, so that the new groups form good interactions with the target surface.

5. The cyclic peptide cP226 (SEQ ID NO: 10) is an example of a compound which binds to the ligand binding site of phospholamban. The structure of cP226 (SEQ

ID NO: 10) can be used as a model for designing new compounds with affinity to phospholamban. Any of the well defined methods for designing peptidomimetics or, more generally, peptide mimics can be used to design such compounds.

A limited number of compounds can be selected through the process outlined above. Anyone skilled with the art would be able to identify such compounds by using the three dimensional structure of phospholamban stored in a computer system. The compounds can be then synthesized and tested for their ability to deactivate phospholamban in an assay similar to that outlined in example 3.

TABLE I. Orthogonal three dimensional coordinates in Ångströms for the cyclic peptide cP226 (SEQ ID NO: 10)

Residue	Atom	X	Y	Z
1	CYS N	-4.500	-5.816	1.065
1	CYS CA	-3.913	-5.081	2.195
1	CYS HN1	-4.805	-6.727	1.378
1	CYS HN2	-5.291	-5.300	0.706
1	CYS HN3	-3.811	-5.926	0.337
1	CYS HA	-4.673	-5.015	2.974
1	CYS C	-3.521	-3.665	1.797
1	CYS O	-4.275	-2.984	1.106
1	CYS CB	-2.716	-5.833	2.773
1	CYS SG	-1.186	-5.807	1.793
1	CYS HB1	-2.485	-5.381	3.737
1	CYS HB2	-3.005	-6.870	2.943
1	CYS LG1	-0.868	-6.293	2.143
1	CYS LG2	-1.298	-5.206	1.500
2	TYR N	-2.334	-3.230	2.226
2	TYR CA	-1.822	-1.920	1.877
2	TYR HN	-1.741	-3.845	2.764
2	TYR HA	-2.467	-1.466	1.124
2	TYR C	-0.427	-2.065	1.301
2	TYR O	-0.005	-3.170	0.967
2	TYR CB	-1.782	-1.016	3.105
2	TYR HB1	-0.796	-0.552	3.147
2	TYR HB2	-2.524	-0.231	2.974
2	TYR CG	-2.052	-1.724	4.413
2	TYR CD1	-3.368	-1.863	4.872
2	TYR HD1	-4.192	-1.509	4.270
2	TYR CD2	-0.987	-2.180	5.200
2	TYR HD2	0.031	-2.071	4.855
2	TYR CE1	-3.619	-2.453	6.116
2	TYR HE1	-4.634	-2.555	6.471

2	TYR	CE2	-1.238	-2.769	6.445
2	TYR	HE2	-0.415	-3.115	7.053
2	TYR	CZ	-2.553	-2.901	6.905
2	TYR	OH	-2.798	-3.464	8.124
2	TYR	HH	-1.998	-3.726	8.585
3	TRP	N	0.295	-0.949	1.195
3	TRP	CA	1.639	-0.990	0.666
3	TRP	HN	-0.084	-0.061	1.491
3	TRP	HA	2.012	-2.004	0.800
3	TRP	C	2.542	-0.026	1.422
3	TRP	O	2.112	0.623	2.372
3	TRP	CB	1.614	-0.654	-0.821
3	TRP	HB1	0.826	-1.238	-1.296
3	TRP	HB2	1.378	0.404	-0.929
3	TRP	CG	2.892	-0.918	-1.546
3	TRP	CD1	3.831	0.008	-1.830
3	TRP	CD2	3.402	-2.178	-2.077
3	TRP	NE1	4.877	-0.580	-2.512
3	TRP	CE2	4.664	-1.934	-2.681
3	TRP	HD1	3.772	1.053	-1.559
3	TRP	HE1	5.708	-0.104	-2.833
3	TRP	CE3	2.933	-3.504	-2.103
3	TRP	HE3	1.978	-3.737	-1.655
3	TRP	CZ2	5.421	-2.947	-3.276
3	TRP	HZ2	6.378	-2.724	-3.723
3	TRP	CZ3	3.684	-4.528	-2.696
3	TRP	HZ3	3.302	-5.538	-2.700
3	TRP	CH2	4.927	-4.255	-3.280
3	TRP	HH2	5.501	-5.051	-3.731
4	GLU	N	3.799	0.056	0.982
4	GLU	CA	4.812	0.899	1.576
4	GLU	HN	4.087	-0.499	0.192
4	GLU	HA	5.405	0.313	2.216
4	GLU	C	4.236	2.017	2.406
4	GLU	O	4.787	2.431	3.423
4	GLU	CB	5.698	1.448	0.481

4	GLU	CG	6.679	0.389	-0.014
4	GLU	CD	7.595	0.956	-1.091
4	GLU	OE1	7.249	0.790	-2.281
4	GLU	OE2	8.627	1.545	-0.704
4	GLU	HB1	5.027	1.763	-0.302
4	GLU	HB2	6.242	2.305	0.856
4	GLU	HG1	7.290	0.056	0.824
4	GLU	HG2	6.128	-0.462	-0.411
5	LEU	N	3.135	2.522	1.889
5	LEU	CA	2.622	3.813	2.150
5	LEU	HN	2.818	2.140	1.031
5	LEU	HA	3.362	4.435	2.574
5	LEU	C	1.437	3.723	3.079
5	LEU	O	1.552	3.459	4.275
5	LEU	CB	2.300	4.379	0.776
5	LEU	HB1	1.233	4.331	0.644
5	LEU	HB2	2.784	3.725	0.059
5	LEU	CG	2.893	5.759	0.516
5	LEU	HG	3.808	5.818	1.092
5	LEU	CD1	3.275	5.906	-0.959
5	LEU	HD11	4.091	5.223	-1.199
5	LEU	HD12	2.426	5.670	-1.594
5	LEU	HD13	3.598	6.928	-1.157
5	LEU	CD2	1.924	6.871	0.916
5	LEU	HD21	1.572	6.707	1.933
5	LEU	HD22	2.430	7.834	0.854
5	LEU	HD23	1.076	6.877	0.232
6	GLU	N	0.310	4.034	2.475
6	GLU	CA	-0.797	4.584	3.234
6	GLU	HN	0.545	4.337	1.543
6	GLU	HA	-0.766	4.194	4.238
6	GLU	C	-2.174	4.334	2.625
6	GLU	O	-3.182	4.404	3.326
6	GLU	CB	-0.517	6.066	3.324
6	GLU	CG	-1.394	6.792	4.342
6	GLU	CD	-1.241	6.210	5.741

6	GLU	OE1	-2.246	5.654	6.235
6	GLU	OE2	-0.124	6.332	6.290
6	GLU	HB1	0.541	6.185	3.551
6	GLU	HB2	-0.707	6.444	2.332
6	GLU	HG1	-1.101	7.841	4.360
6	GLU	HG2	-2.436	6.722	4.036
7	TRP	N	-2.229	4.100	1.315
7	TRP	CA	-3.460	4.217	0.555
7	TRP	HN	-1.368	3.977	0.804
7	TRP	HA	-4.317	4.129	1.222
7	TRP	C	-3.504	3.121	-0.491
7	TRP	O	-4.454	2.344	-0.558
7	TRP	CB	-3.485	5.577	-0.150
7	TRP	HB1	-4.435	5.670	-0.675
7	TRP	HB2	-3.431	6.359	0.607
7	TRP	CG	-2.379	5.797	-1.141
7	TRP	CD1	-1.059	5.740	-0.857
7	TRP	CD2	-2.452	6.056	-2.579
7	TRP	NE1	-0.316	5.931	-1.995
7	TRP	CE2	-1.127	6.116	-3.086
7	TRP	HD1	-0.629	5.529	0.104
7	TRP	HE1	0.688	5.926	-2.054
7	TRP	CE3	-3.486	6.229	-3.517
7	TRP	HE3	-4.514	6.191	-3.189
7	TRP	CZ2	-0.832	6.315	-4.430
7	TRP	HZ2	0.208	6.311	-4.723
7	TRP	CZ3	-3.203	6.453	-4.874
7	TRP	HZ3	-4.014	6.590	-5.573
7	TRP	CH2	-1.880	6.491	-5.336
7	TRP	HH2	-1.673	6.653	-6.384
8	LEU	N	-2.458	3.080	-1.316
8	LEU	CA	-2.399	2.215	-2.463
8	LEU	HN	-1.741	3.788	-1.251
8	LEU	HA	-3.342	2.268	-2.987
8	LEU	C	-2.138	0.775	-2.047
8	LEU	O	-2.050	0.466	-0.861

8	LEU	CB	-1.377	2.801	-3.420
8	LEU	HB1	-1.919	3.553	-3.985
8	LEU	HB2	-0.627	3.313	-2.838
8	LEU	CG	-0.697	1.810	-4.364
8	LEU	HG	-1.411	1.066	-4.715
8	LEU	CD1	0.486	1.139	-3.660
8	LEU	HD11	0.148	0.611	-2.770
8	LEU	HD12	0.958	0.428	-4.336
8	LEU	HD13	1.216	1.894	-3.367
8	LEU	CD2	-0.166	2.597	-5.559
8	LEU	HD21	0.377	1.929	-6.226
8	LEU	HD22	-0.999	3.047	-6.099
8	LEU	HD23	0.503	3.385	-5.211
9	PRO	N	-2.105	-0.108	-3.045
9	PRO	CA	-2.388	-1.520	-2.908
9	PRO	CD	-2.238	0.281	-4.432
9	PRO	HA	-3.411	-1.651	-2.592
9	PRO	HD1	-1.244	0.463	-4.841
9	PRO	HD2	-2.849	1.180	-4.534
9	PRO	C	-1.489	-2.327	-2.003
9	PRO	O	-0.712	-1.817	-1.200
9	PRO	CB	-2.321	-2.090	-4.308
9	PRO	HB1	-2.906	-3.003	-4.418
9	PRO	HB2	-1.278	-2.261	-4.524
9	PRO	CG	-2.874	-0.932	-5.115
9	PRO	HG1	-3.943	-0.968	-4.924
9	PRO	HG2	-2.647	-0.998	-6.179
10	CYS	N	-1.663	-3.631	-2.196
10	CYS	CA	-1.012	-4.692	-1.465
10	CYS	HN	-2.332	-3.890	-2.912
10	CYS	HA	-0.585	-4.387	-0.542
10	CYS	C	0.010	-5.351	-2.349
10	CYS	O	1.190	-5.013	-2.406
10	CYS	CB	-2.057	-5.730	-1.120
10	CYS	SG	-1.557	-6.937	0.133
10	CYS	HB1	-2.248	-6.248	-2.047

10	CYS	HB2	-2.961	-5.214	-0.820
10	CYS	LG1	-2.185	-7.126	0.309
10	CYS	LG2	-0.911	-6.990	-0.068
11	ALA	N	-0.563	-6.322	-3.032
11	ALA	CA	0.071	-7.137	-4.047
11	ALA	HN	-1.541	-6.400	-2.791
11	ALA	HA	-0.700	-7.659	-4.614
11	ALA	C	0.986	-8.164	-3.395
11	ALA	O	1.316	-7.956	-2.208
11	ALA	OXT	1.336	-9.138	-4.095
11	ALA	CB	0.852	-6.232	-4.994
11	ALA	HB1	1.630	-5.713	-4.434
11	ALA	HB2	1.304	-6.831	-5.783
11	ALA	HB3	0.172	-5.502	-5.433

END

TABLE III. Orthogonal three dimensional coordinates in Ångströms for the phospholamban (1-36) peptide (SEQ ID NO:9)

Residue	Atom	X	Y	Z
1	MET N	-18.883	-3.493	9.020
1	MET CA	-17.518	-3.957	8.653
1	MET HN1	-19.536	-4.279	9.119
1	MET HN2	-18.890	-2.989	9.914
1	MET HN3	-19.280	-2.863	8.314
1	MET HA	-16.901	-3.038	8.626
1	MET C	-17.415	-4.543	7.212
1	MET O	-16.572	-4.081	6.439
1	MET CB	-16.875	-4.826	9.772
1	MET HB1	-15.795	-4.924	9.547
1	MET HB2	-16.893	-4.265	10.726
1	MET CG	-17.422	-6.248	10.031
1	MET SD	-19.151	-6.241	10.562
1	MET CE	-18.991	-5.887	12.325
1	MET HG1	-17.330	-6.874	9.124
1	MET HG2	-16.810	-6.758	10.798
1	MET HE1	-19.984	-5.878	12.809
1	MET HE2	-18.374	-6.655	12.827
1	MET HE3	-18.521	-4.900	12.492
2	GLU N	-18.254	-5.533	6.842
2	GLU CA	-18.265	-6.128	5.478
2	GLU HN	-18.900	-5.856	7.571
2	GLU HA	-17.210	-6.282	5.173
2	GLU C	-18.948	-5.225	4.401
2	GLU O	-19.754	-4.342	4.720
2	GLU CB	-18.906	-7.547	5.541
2	GLU CG	-20.416	-7.627	5.896
2	GLU CD	-21.035	-8.987	5.590
2	GLU OE1	-21.495	-9.191	4.444

2	GLU	OE2	-21.070	-9.854	6.489
2	GLU	HB1	-18.746	-8.031	4.557
2	GLU	HB2	-18.337	-8.184	6.246
2	GLU	HG1	-20.571	-7.388	6.963
2	GLU	HG2	-20.990	-6.864	5.338
3	LYS	N	-18.664	-5.521	3.116
3	LYS	CA	-19.464	-5.049	1.948
3	LYS	HN	-17.993	-6.291	3.013
3	LYS	HA	-19.008	-5.581	1.092
3	LYS	C	-19.248	-3.539	1.622
3	LYS	O	-18.370	-3.219	0.818
3	LYS	CB	-20.930	-5.587	1.997
3	LYS	CG	-21.709	-5.508	0.666
3	LYS	CD	-23.177	-5.990	0.736
3	LYS	CE	-23.411	-7.513	0.856
3	LYS	NZ	-23.320	-8.021	2.243
3	LYS	HB1	-20.905	-6.644	2.322
3	LYS	HB2	-21.495	-5.058	2.788
3	LYS	HG1	-21.711	-4.458	0.320
3	LYS	HG2	-21.169	-6.068	-0.122
3	LYS	HD1	-23.726	-5.441	1.525
3	LYS	HD2	-23.662	-5.662	-0.204
3	LYS	HE1	-24.420	-7.750	0.465
3	LYS	HE2	-22.711	-8.069	0.201
3	LYS	HZ1	-22.376	-7.928	2.639
3	LYS	HZ2	-23.954	-7.523	2.880
3	LYS	HZ3	-23.558	-9.017	2.311
4	VAL	N	-20.017	-2.625	2.249
4	VAL	CA	-19.844	-1.150	2.084
4	VAL	HN	-20.672	-3.020	2.932
4	VAL	HA	-19.723	-0.979	0.995
4	VAL	C	-18.532	-0.595	2.734
4	VAL	O	-17.763	0.075	2.042
4	VAL	CB	-21.166	-0.392	2.460

4	VAL	HB	-21.981	-0.865	1.880
4	VAL	CG1	-21.148	1.087	2.010
4	VAL	HG11	-22.125	1.579	2.173
4	VAL	HG12	-20.389	1.679	2.555
4	VAL	HG13	-20.921	1.182	0.931
4	VAL	CG2	-21.594	-0.469	3.946
4	VAL	HG21	-22.587	-0.009	4.106
4	VAL	HG22	-21.672	-1.514	4.297
4	VAL	HG23	-20.886	0.051	4.615
5	GLN	N	-18.257	-0.895	4.022
5	GLN	CA	-16.969	-0.532	4.685
5	GLN	HN	-18.970	-1.467	4.487
5	GLN	HA	-16.753	0.525	4.431
5	GLN	C	-15.708	-1.326	4.214
5	GLN	O	-14.609	-0.764	4.263
5	GLN	CB	-17.126	-0.595	6.230
5	GLN	HB1	-16.134	-0.491	6.711
5	GLN	HB2	-17.470	-1.605	6.519
5	GLN	CG	-18.072	0.452	6.870
5	GLN	HG1	-19.093	0.350	6.454
5	GLN	HG2	-18.189	0.209	7.943
5	GLN	CD	-17.577	1.906	6.777
5	GLN	OE1	-16.699	2.331	7.525
5	GLN	NE2	-18.118	2.697	5.865
5	GLN	HE21	-17.754	3.655	5.830
5	GLN	HE22	-18.828	2.281	5.254
6	TYR	N	-15.837	-2.578	3.722
6	TYR	CA	-14.754	-3.255	2.953
6	TYR	HN	-16.798	-2.934	3.726
6	TYR	HA	-13.836	-3.179	3.569
6	TYR	C	-14.429	-2.578	1.577
6	TYR	O	-13.246	-2.405	1.284
6	TYR	CB	-15.073	-4.771	2.823
6	TYR	HB1	-15.450	-5.161	3.789

6	TYR	HB2	-15.922	-4.917	2.126
6	TYR	CG	-13.870	-5.631	2.391
6	TYR	CD1	-12.947	-6.081	3.342
6	TYR	HD1	-13.093	-5.867	4.391
6	TYR	CD2	-13.656	-5.919	1.038
6	TYR	HD2	-14.351	-5.571	0.287
6	TYR	CE1	-11.820	-6.795	2.944
6	TYR	HE1	-11.109	-7.135	3.683
6	TYR	CE2	-12.526	-6.633	0.642
6	TYR	HE2	-12.364	-6.840	-0.406
6	TYR	CZ	-11.608	-7.066	1.594
6	TYR	OH	-10.486	-7.749	1.203
6	TYR	HH	-10.543	-7.928	0.261
7	LEU	N	-15.435	-2.177	0.768
7	LEU	CA	-15.228	-1.333	-0.448
7	LEU	HN	-16.377	-2.392	1.118
7	LEU	HA	-14.505	-1.879	-1.083
7	LEU	C	-14.618	0.088	-0.187
7	LEU	O	-13.755	0.514	-0.959
7	LEU	CB	-16.568	-1.276	-1.239
7	LEU	HB1	-16.964	-2.304	-1.362
7	LEU	HB2	-17.320	-0.763	-0.607
7	LEU	CG	-16.545	-0.600	-2.641
7	LEU	HG	-16.134	0.421	-2.532
7	LEU	CD1	-17.977	-0.449	-3.189
7	LEU	HD11	-18.471	-1.427	-3.341
7	LEU	HD12	-18.616	0.136	-2.501
7	LEU	HD13	-17.989	0.081	-4.159
7	LEU	CD2	-15.670	-1.352	-3.665
7	LEU	HD21	-16.014	-2.391	-3.823
7	LEU	HD22	-15.676	-0.852	-4.652
7	LEU	HD23	-14.613	-1.398	-3.347
8	THR	N	-15.020	0.793	0.895
8	THR	CA	-14.337	2.035	1.379

8	THR	HN	-15.786	0.362	1.425
8	THR	HA	-14.401	2.775	0.558
8	THR	C	-12.813	1.822	1.687
8	THR	O	-11.983	2.555	1.144
8	THR	CB	-15.125	2.633	2.591
8	THR	OG1	-16.503	2.806	2.270
8	THR	HB	-15.061	1.931	3.445
8	THR	HG1	-16.535	3.486	1.594
8	THR	CG2	-14.620	4.006	3.070
8	THR	HG21	-14.652	4.764	2.265
8	THR	HG22	-15.233	4.389	3.908
8	THR	HG23	-13.579	3.955	3.436
9	ARG	N	-12.454	0.800	2.491
9	ARG	CA	-11.040	0.367	2.694
9	ARG	HN	-13.242	0.262	2.868
9	ARG	HA	-10.475	1.248	3.056
9	ARG	C	-10.279	-0.141	1.422
9	ARG	O	-9.068	0.074	1.342
9	ARG	CB	-10.986	-0.704	3.816
9	ARG	CG	-11.336	-0.195	5.239
9	ARG	CD	-11.278	-1.278	6.331
9	ARG	NE	-12.399	-2.249	6.225
9	ARG	CZ	-12.453	-3.428	6.870
9	ARG	NH1	-13.518	-4.178	6.704
9	ARG	NH2	-11.493	-3.881	7.666
9	ARG	HB1	-9.966	-1.132	3.864
9	ARG	HB2	-11.641	-1.555	3.549
9	ARG	HG1	-10.627	0.611	5.509
9	ARG	HG2	-12.331	0.292	5.247
9	ARG	HD1	-10.296	-1.790	6.290
9	ARG	HD2	-11.320	-0.790	7.325
9	ARG	HE	-13.225	-2.039	5.652
9	ARG	HH12	-14.259	-3.806	6.101
9	ARG	HH11	-13.535	-5.071	7.209

9	ARG	HH21	-10.677	-3.270	7.781
9	ARG	HH22	-11.642	-4.793	8.108
10	SER	N	-10.946	-0.787	0.443
10	SER	CA	-10.354	-1.132	-0.884
10	SER	HN	-11.942	-0.946	0.641
10	SER	HA	-9.439	-1.724	-0.697
10	SER	C	-9.924	0.093	-1.755
10	SER	O	-8.816	0.081	-2.300
10	SER	CB	-11.306	-2.070	-1.661
10	SER	OG	-11.499	-3.303	-0.971
10	SER	HB1	-12.283	-1.585	-1.840
10	SER	HB2	-10.895	-2.294	-2.663
10	SER	HG	-12.151	-3.796	-1.475
11	ALA	N	-10.757	1.151	-1.848
11	ALA	CA	-10.344	2.469	-2.403
11	ALA	HN	-11.651	1.031	-1.357
11	ALA	HA	-9.966	2.296	-3.429
11	ALA	C	-9.219	3.223	-1.619
11	ALA	O	-8.321	3.776	-2.259
11	ALA	CB	-11.608	3.341	-2.539
11	ALA	HB1	-12.075	3.558	-1.559
11	ALA	HB2	-11.379	4.314	-3.012
11	ALA	HB3	-12.378	2.857	-3.170
12	ILE	N	-9.242	3.226	-0.266
12	ILE	CA	-8.172	3.838	0.586
12	ILE	HN	-10.056	2.754	0.145
12	ILE	HA	-8.006	4.865	0.200
12	ILE	C	-6.795	3.099	0.438
12	ILE	O	-5.794	3.779	0.207
12	ILE	CB	-8.658	4.011	2.076
12	ILE	HB	-8.948	3.006	2.440
12	ILE	CG1	-9.901	4.950	2.214
12	ILE	HG11	-10.620	4.757	1.398
12	ILE	HG12	-9.602	6.005	2.066

12	ILE	CG2	-7.544	4.533	3.028
12	ILE	HG21	-7.887	4.618	4.075
12	ILE	HG22	-6.669	3.857	3.055
12	ILE	HG23	-7.174	5.529	2.721
12	ILE	CD1	-10.686	4.819	3.530
12	ILE	HD11	-11.582	5.467	3.523
12	ILE	HD12	-11.037	3.782	3.694
12	ILE	HD13	-10.086	5.112	4.410
13	ARG	N	-6.728	1.753	0.563
13	ARG	CA	-5.468	0.976	0.354
13	ARG	HN	-7.628	1.294	0.749
13	ARG	HA	-4.726	1.433	1.041
13	ARG	C	-4.830	1.090	-1.073
13	ARG	O	-3.613	1.260	-1.173
13	ARG	CB	-5.630	-0.486	0.871
13	ARG	CG	-6.342	-1.498	-0.063
13	ARG	CD	-6.640	-2.883	0.549
13	ARG	NE	-7.896	-2.883	1.345
13	ARG	CZ	-8.627	-3.976	1.626
13	ARG	NH1	-8.236	-5.216	1.363
13	ARG	NH2	-9.800	-3.810	2.198
13	ARG	HB1	-6.139	-0.472	1.855
13	ARG	HB2	-4.622	-0.886	1.095
13	ARG	HG1	-7.266	-1.050	-0.471
13	ARG	HG2	-5.699	-1.654	-0.950
13	ARG	HD1	-6.722	-3.609	-0.283
13	ARG	HD2	-5.789	-3.231	1.168
13	ARG	HE	-8.340	-1.998	1.613
13	ARG	HH12	-7.314	-5.315	0.927
13	ARG	HH11	-8.879	-5.974	1.620
13	ARG	HH21	-10.096	-2.844	2.364
13	ARG	HH22	-10.347	-4.658	2.382
14	ARG	N	-5.647	1.040	-2.150
14	ARG	CA	-5.205	1.316	-3.547

14 ARG	HN	-6.640	0.917	-1.918
14 ARG	HA	-4.382	0.609	-3.774
14 ARG	C	-4.654	2.762	-3.783
14 ARG	O	-3.574	2.899	-4.362
14 ARG	CB	-6.387	0.954	-4.492
14 ARG	CG	-6.046	0.936	-6.002
14 ARG	CD	-7.241	0.641	-6.935
14 ARG	NE	-7.781	-0.743	-6.818
14 ARG	CZ	-7.326	-1.813	-7.495
14 ARG	NH1	-6.304	-1.776	-8.341
14 ARG	NH2	-7.927	-2.967	-7.304
14 ARG	HB1	-6.779	-0.047	-4.223
14 ARG	HB2	-7.232	1.648	-4.313
14 ARG	HG1	-5.635	1.922	-6.287
14 ARG	HG2	-5.227	0.217	-6.196
14 ARG	HD1	-8.058	1.355	-6.720
14 ARG	HD2	-6.959	0.862	-7.983
14 ARG	HE	-8.576	-0.946	-6.202
14 ARG	HH12	-5.856	-0.864	-8.466
14 ARG	HH11	-6.046	-2.656	-8.800
14 ARG	HH21	-8.716	-2.973	-6.649
14 ARG	HH22	-7.566	-3.769	-7.830
15 ALA	N	-5.366	3.815	-3.329
15 ALA	CA	-4.859	5.215	-3.352
15 ALA	HN	-6.249	3.572	-2.865
15 ALA	HA	-4.610	5.460	-4.404
15 ALA	C	-3.580	5.502	-2.503
15 ALA	O	-2.687	6.192	-2.996
15 ALA	CB	-6.019	6.148	-2.948
15 ALA	HB1	-5.723	7.211	-3.012
15 ALA	HB2	-6.898	6.024	-3.608
15 ALA	HB3	-6.360	5.965	-1.911
16 SER	N	-3.471	4.962	-1.270
16 SER	CA	-2.253	5.080	-0.420

16	SER	HN	-4.293	4.425	-0.965
16	SER	HA	-2.050	6.162	-0.292
16	SER	C	-0.958	4.438	-1.011
16	SER	O	0.081	5.105	-1.033
16	SER	CB	-2.595	4.527	0.984
16	SER	OG	-1.533	4.766	1.904
16	SER	HB1	-2.808	3.441	0.942
16	SER	HB2	-3.513	5.000	1.382
16	SER	HG	-0.749	4.360	1.526
17	THR	N	-1.011	3.178	-1.503
17	THR	CA	0.124	2.551	-2.246
17	THR	HN	-1.934	2.731	-1.438
17	THR	HA	1.023	2.737	-1.625
17	THR	C	0.453	3.189	-3.638
17	THR	O	1.637	3.295	-3.969
17	THR	CB	-0.014	0.998	-2.268
17	THR	OG1	1.246	0.419	-2.593
17	THR	HB	-0.270	0.656	-1.245
17	THR	HG1	1.437	0.690	-3.494
17	THR	CG2	-1.058	0.403	-3.230
17	THR	HG21	-1.179	-0.683	-3.070
17	THR	HG22	-2.050	0.867	-3.099
17	THR	HG23	-0.770	0.546	-4.287
18	ILE	N	-0.556	3.629	-4.427
18	ILE	CA	-0.333	4.400	-5.692
18	ILE	HN	-1.497	3.496	-4.036
18	ILE	HA	0.698	4.211	-6.053
18	ILE	C	-0.421	5.946	-5.379
18	ILE	O	-1.236	6.672	-5.957
18	ILE	CB	-1.288	3.908	-6.852
18	ILE	HB	-2.305	4.271	-6.603
18	ILE	CG1	-1.395	2.357	-7.037
18	ILE	HG11	-1.526	1.871	-6.054
18	ILE	HG12	-0.444	1.948	-7.427

18 ILE	CG2	-0.883	4.538	-8.215
18 ILE	HG21	-0.844	5.641	-8.182
18 ILE	HG22	0.115	4.193	-8.547
18 ILE	HG23	-1.597	4.290	-9.020
18 ILE	CD1	-2.570	1.867	-7.901
18 ILE	HD11	-2.470	2.172	-8.958
18 ILE	HD12	-2.641	0.764	-7.888
18 ILE	HD13	-3.536	2.260	-7.531
19 GLU	N	0.446	6.441	-4.470
19 GLU	CA	0.634	7.888	-4.165
19 GLU	HN	1.033	5.726	-4.025
19 GLU	HA	0.850	8.416	-5.116
19 GLU	C	1.890	7.939	-3.246
19 GLU	O	1.773	7.912	-2.015
19 GLU	CB	-0.615	8.561	-3.518
19 GLU	CG	-0.502	10.091	-3.331
19 GLU	CD	-1.757	10.709	-2.714
19 GLU	OE1	-1.821	10.833	-1.471
19 GLU	OE2	-2.681	11.079	-3.471
19 GLU	HB1	-1.509	8.364	-4.140
19 GLU	HB2	-0.836	8.082	-2.544
19 GLU	HG1	0.364	10.339	-2.690
19 GLU	HG2	-0.306	10.582	-4.302
20 MET	N	3.097	7.953	-3.849
20 MET	CA	4.367	7.766	-3.099
20 MET	HN	3.066	7.947	-4.874
20 MET	HA	4.229	8.109	-2.058
20 MET	C	5.500	8.612	-3.764
20 MET	O	5.912	8.250	-4.873
20 MET	CB	4.774	6.264	-3.011
20 MET	HB1	4.730	5.784	-4.006
20 MET	HB2	5.837	6.193	-2.714
20 MET	CG	3.968	5.437	-1.993
20 MET	SD	4.791	3.860	-1.711

20 MET	CE	4.009	3.366	-0.164
20 MET	HG1	3.884	5.980	-1.034
20 MET	HG2	2.935	5.259	-2.347
20 MET	HE1	4.407	2.394	0.179
20 MET	HE2	4.198	4.113	0.628
20 MET	HE3	2.916	3.266	-0.289
21 PRO	N	6.082	9.682	-3.136
21 PRO	CA	7.307	10.353	-3.661
21 PRO	CD	5.577	10.289	-1.885
21 PRO	HA	7.190	10.566	-4.742
21 PRO	HD1	5.491	9.562	-1.057
21 PRO	HD2	4.577	10.737	-2.051
21 PRO	C	8.610	9.505	-3.466
21 PRO	O	8.564	8.355	-3.015
21 PRO	CB	7.264	11.689	-2.884
21 PRO	HB1	8.254	12.168	-2.762
21 PRO	HB2	6.636	12.421	-3.431
21 PRO	CG	6.612	11.360	-1.542
21 PRO	HG1	6.162	12.247	-1.057
21 PRO	HG2	7.367	10.953	-0.842
22 GLN	N	9.775	10.079	-3.825
22 GLN	CA	11.099	9.383	-3.763
22 GLN	HN	9.684	11.028	-4.202
22 GLN	HA	11.033	8.583	-4.525
22 GLN	C	11.482	8.638	-2.439
22 GLN	O	11.949	7.500	-2.513
22 GLN	CB	12.223	10.322	-4.282
22 GLN	HB1	13.148	9.727	-4.407
22 GLN	HB2	11.966	10.640	-5.311
22 GLN	CG	12.550	11.578	-3.435
22 GLN	HG1	11.630	12.168	-3.268
22 GLN	HG2	12.884	11.281	-2.422
22 GLN	CD	13.588	12.498	-4.093
22 GLN	OE1	13.260	13.333	-4.933

22 GLN NE2	14.855	12.373	-3.736
22 GLN HE21	15.067	11.660	-3.030
22 GLN HE22	15.516	13.007	-4.196
23 GLN N	11.241	9.236	-1.255
23 GLN CA	11.413	8.552	0.061
23 GLN HN	10.841	10.178	-1.328
23 GLN HA	12.430	8.113	0.051
23 GLN C	10.448	7.350	0.351
23 GLN O	10.910	6.324	0.858
23 GLN CB	11.447	9.600	1.208
23 GLN HB1	12.280	10.306	1.013
23 GLN HB2	11.743	9.086	2.143
23 GLN CG	10.154	10.415	1.475
23 GLN HG1	9.839	10.931	0.549
23 GLN HG2	9.316	9.735	1.719
23 GLN CD	10.320	11.475	2.574
23 GLN OE1	10.799	12.580	2.329
23 GLN NE2	9.928	11.173	3.800
23 GLN HE21	9.539	10.235	3.942
23 GLN HE22	10.049	11.909	4.503
24 ALA N	9.143	7.463	0.030
24 ALA CA	8.159	6.359	0.204
24 ALA HN	8.886	8.352	-0.413
24 ALA HA	8.289	5.956	1.227
24 ALA C	8.290	5.147	-0.774
24 ALA O	8.102	4.010	-0.329
24 ALA CB	6.748	6.969	0.163
24 ALA HB1	6.615	7.761	0.924
24 ALA HB2	6.522	7.421	-0.821
24 ALA HB3	5.969	6.209	0.360
25 ARG N	8.632	5.358	-2.064
25 ARG CA	9.043	4.252	-2.981
25 ARG HN	8.743	6.347	-2.322
25 ARG HA	8.300	3.441	-2.842

25 ARG C	10.430	3.587	-2.668
25 ARG O	10.540	2.365	-2.801
25 ARG CB	8.889	4.651	-4.473
25 ARG CG	9.786	5.799	-5.012
25 ARG CD	10.177	5.685	-6.502
25 ARG NE	11.091	4.541	-6.782
25 ARG CZ	12.421	4.541	-6.578
25 ARG NH1	13.108	5.603	-6.180
25 ARG NH2	13.079	3.421	-6.783
25 ARG HB1	7.831	4.908	-4.677
25 ARG HB2	9.054	3.736	-5.072
25 ARG HG1	9.259	6.756	-4.857
25 ARG HG2	10.708	5.894	-4.408
25 ARG HD1	9.264	5.572	-7.115
25 ARG HD2	10.627	6.636	-6.847
25 ARG HE	10.720	3.637	-7.092
25 ARG HH12	12.567	6.460	-6.027
25 ARG HH11	14.116	5.475	-6.041
25 ARG HH21	12.529	2.612	-7.087
25 ARG HH22	14.087	3.441	-6.595
26 GLN N	11.453	4.350	-2.215
26 GLN CA	12.697	3.779	-1.612
26 GLN HN	11.240	5.353	-2.152
26 GLN HA	13.138	3.102	-2.365
26 GLN C	12.484	2.907	-0.334
26 GLN O	13.148	1.878	-0.206
26 GLN CB	13.749	4.894	-1.344
26 GLN HB1	13.273	5.724	-0.787
26 GLN HB2	14.530	4.520	-0.653
26 GLN CG	14.472	5.467	-2.589
26 GLN HG1	15.030	6.376	-2.295
26 GLN HG2	13.724	5.828	-3.317
26 GLN CD	15.422	4.491	-3.305
26 GLN OE1	15.071	3.875	-4.309

26 GLN	NE2	16.638	4.327	-2.815
26 GLN	HE21	16.862	4.843	-1.957
26 GLN	HE22	17.228	3.639	-3.294
27 LYS	N	11.548	3.264	0.572
27 LYS	CA	11.074	2.367	1.667
27 LYS	HN	11.088	4.160	0.371
27 LYS	HA	11.940	2.194	2.336
27 LYS	C	10.559	0.961	1.203
27 LYS	O	10.956	-0.041	1.801
27 LYS	CB	10.012	3.154	2.484
27 LYS	CG	9.566	2.501	3.813
27 LYS	CD	8.426	3.241	4.552
27 LYS	CE	6.988	2.966	4.055
27 LYS	NZ	6.636	3.687	2.812
27 LYS	HB1	9.132	3.324	1.842
27 LYS	HB2	10.393	4.168	2.720
27 LYS	HG1	9.272	1.447	3.652
27 LYS	HG2	10.446	2.449	4.483
27 LYS	HD1	8.466	2.914	5.609
27 LYS	HD2	8.630	4.328	4.599
27 LYS	HE1	6.828	1.878	3.918
27 LYS	HE2	6.273	3.264	4.847
27 LYS	HZ1	6.751	4.702	2.914
27 LYS	HZ2	7.225	3.398	2.022
27 LYS	HZ3	5.663	3.522	2.532
28 LEU	N	9.719	0.878	0.145
28 LEU	CA	9.260	-0.418	-0.438
28 LEU	HN	9.490	1.779	-0.290
28 LEU	HA	8.796	-0.997	0.384
28 LEU	C	10.403	-1.321	-1.004
28 LEU	O	10.505	-2.475	-0.581
28 LEU	CB	8.156	-0.187	-1.513
28 LEU	HB1	8.584	0.438	-2.322
28 LEU	HB2	7.934	-1.155	-2.004

28 LEU	CG	6.803	0.435	-1.064
28 LEU	HG	7.002	1.418	-0.596
28 LEU	CD1	5.909	0.683	-2.295
28 LEU	HD11	5.692	-0.251	-2.848
28 LEU	HD12	4.935	1.125	-2.018
28 LEU	HD13	6.384	1.382	-3.009
28 LEU	CD2	6.044	-0.436	-0.042
28 LEU	HD21	5.068	0.007	0.230
28 LEU	HD22	5.844	-1.452	-0.431
28 LEU	HD23	6.607	-0.550	0.901
29 GLN	N	11.255	-0.816	-1.925
29 GLN	CA	12.406	-1.595	-2.475
29 GLN	HN	11.082	0.162	-2.182
29 GLN	HA	11.991	-2.584	-2.755
29 GLN	C	13.567	-1.934	-1.484
29 GLN	O	14.147	-3.014	-1.607
29 GLN	CB	12.883	-1.003	-3.833
29 GLN	HB1	13.599	-1.706	-4.301
29 GLN	HB2	12.019	-1.021	-4.524
29 GLN	CG	13.475	0.432	-3.883
29 GLN	HG1	12.803	1.117	-3.337
29 GLN	HG2	13.438	0.795	-4.926
29 GLN	CD	14.910	0.617	-3.360
29 GLN	OE1	15.141	1.212	-2.310
29 GLN	NE2	15.908	0.140	-4.084
29 GLN	HE21	15.657	-0.353	-4.946
29 GLN	HE22	16.847	0.271	-3.695
30 ASN	N	13.883	-1.064	-0.500
30 ASN	CA	14.859	-1.363	0.589
30 ASN	HN	13.357	-0.182	-0.525
30 ASN	HA	15.789	-1.718	0.103
30 ASN	C	14.394	-2.487	1.576
30 ASN	O	15.180	-3.393	1.865
30 ASN	CB	15.209	-0.024	1.297

30 ASN	HB1	14.313	0.381	1.811
30 ASN	HB2	15.459	0.741	0.538
30 ASN	CG	16.411	-0.079	2.257
30 ASN	OD1	17.565	-0.101	1.835
30 ASN	ND2	16.175	-0.094	3.557
30 ASN	HD21	15.192	-0.090	3.848
30 ASN	HD22	17.001	-0.122	4.163
31 LEU	N	13.133	-2.449	2.059
31 LEU	CA	12.510	-3.577	2.815
31 LEU	HN	12.579	-1.645	1.742
31 LEU	HA	13.171	-3.801	3.673
31 LEU	C	12.370	-4.919	2.020
31 LEU	O	12.617	-5.979	2.598
31 LEU	CB	11.135	-3.137	3.397
31 LEU	HB1	10.484	-2.835	2.550
31 LEU	HB2	10.622	-4.020	3.828
31 LEU	CG	11.136	-2.012	4.471
31 LEU	HG	11.710	-1.150	4.082
31 LEU	CD1	9.702	-1.513	4.733
31 LEU	HD11	9.051	-2.309	5.141
31 LEU	HD12	9.685	-0.673	5.453
31 LEU	HD13	9.225	-1.143	3.806
31 LEU	CD2	11.790	-2.446	5.800
31 LEU	HD21	11.766	-1.636	6.552
31 LEU	HD22	11.283	-3.323	6.245
31 LEU	HD23	12.853	-2.715	5.665
32 PHE	N	12.006	-4.880	0.720
32 PHE	CA	11.945	-6.085	-0.153
32 PHE	HN	11.808	-3.940	0.359
32 PHE	HA	11.330	-6.832	0.380
32 PHE	C	13.325	-6.763	-0.442
32 PHE	O	13.428	-7.981	-0.273
32 PHE	CB	11.156	-5.718	-1.445
32 PHE	HB1	10.236	-5.162	-1.176

32 PHE	HB2	11.739	-4.989	-2.041
32 PHE	CG	10.748	-6.916	-2.322
32 PHE	CD1	11.529	-7.286	-3.422
32 PHE	HD1	12.424	-6.733	-3.671
32 PHE	CD2	9.607	-7.664	-2.009
32 PHE	HD2	8.994	-7.398	-1.160
32 PHE	CE1	11.181	-8.394	-4.190
32 PHE	HE1	11.800	-8.686	-5.026
32 PHE	CE2	9.262	-8.773	-2.778
32 PHE	HE2	8.387	-9.355	-2.525
32 PHE	CZ	10.049	-9.138	-3.867
32 PHE	HZ	9.787	-10.004	-4.457
33 ILE	N	14.353	-6.004	-0.883
33 ILE	CA	15.663	-6.564	-1.345
33 ILE	HN	14.134	-5.006	-0.980
33 ILE	HA	15.384	-7.289	-2.133
33 ILE	C	16.495	-7.409	-0.317
33 ILE	O	17.181	-8.344	-0.739
33 ILE	CB	16.489	-5.449	-2.086
33 ILE	HB	15.776	-4.903	-2.735
33 ILE	CG1	17.557	-6.044	-3.052
33 ILE	HG11	17.119	-6.899	-3.603
33 ILE	HG12	18.393	-6.481	-2.473
33 ILE	CG2	17.113	-4.397	-1.130
33 ILE	HG21	17.525	-3.530	-1.679
33 ILE	HG22	16.366	-3.996	-0.425
33 ILE	HG23	17.934	-4.823	-0.524
33 ILE	CD1	18.119	-5.067	-4.098
33 ILE	HD11	17.319	-4.642	-4.731
33 ILE	HD12	18.660	-4.224	-3.631
33 ILE	HD13	18.834	-5.575	-4.772
34 ASN	N	16.432	-7.113	1.000
34 ASN	CA	17.066	-7.958	2.057
34 ASN	HN	15.817	-6.323	1.227

34 ASN	HA	18.139	-8.016	1.783
34 ASN	C	16.596	-9.450	2.131
34 ASN	O	17.449	-10.334	2.262
34 ASN	CB	17.063	-7.233	3.433
34 ASN	HB1	17.629	-7.850	4.158
34 ASN	HB2	17.675	-6.313	3.346
34 ASN	CG	15.701	-6.904	4.084
34 ASN	OD1	14.944	-7.786	4.489
34 ASN	ND2	15.366	-5.635	4.221
34 ASN	HD21	16.007	-4.942	3.822
34 ASN	HD22	14.441	-5.462	4.626
35 PHE	N	15.282	-9.732	2.018
35 PHE	CA	14.756	-11.120	1.861
35 PHE	HN	14.679	-8.909	1.899
35 PHE	HA	15.444	-11.796	2.407
35 PHE	C	14.743	-11.669	0.395
35 PHE	O	15.062	-12.848	0.209
35 PHE	CB	13.404	-11.302	2.611
35 PHE	HB1	13.580	-11.112	3.689
35 PHE	HB2	13.142	-12.378	2.592
35 PHE	CG	12.173	-10.489	2.152
35 PHE	CD1	11.351	-10.970	1.126
35 PHE	HD1	11.594	-11.891	0.615
35 PHE	CD2	11.830	-9.300	2.804
35 PHE	HD2	12.449	-8.910	3.599
35 PHE	CE1	10.209	-10.267	0.752
35 PHE	HE1	9.583	-10.638	-0.047
35 PHE	CE2	10.675	-8.610	2.442
35 PHE	HE2	10.412	-7.694	2.952
35 PHE	CZ	9.867	-9.092	1.417
35 PHE	HZ	8.977	-8.551	1.131
36 CYS	N	14.361	-10.862	-0.617
36 CYS	CA	14.287	-11.298	-2.032
36 CYS	C	14.877	-10.182	-2.921

36	CYS	O	16.060	-10.297	-3.312
36	CYS	CB	12.828	-11.666	-2.386
36	CYS	SG	12.713	-12.287	-4.099
36	CYS	OXT	14.177	-9.193	-3.237
36	CYS	HN	14.138	-9.897	-0.344
36	CYS	HA	14.907	-12.202	-2.195
36	CYS	HB1	12.450	-12.455	-1.710
36	CYS	HB2	12.149	-10.800	-2.263
36	CYS	HG	12.819	-11.106	-4.704

END

TABLE IV. Orthogonal three dimensional coordinates in Ångströms for the cP226•phospholamban(1-36) complex

Residue	Atom	X	Y	Z
1	MET N	-18.366	-10.441	5.382
1	MET CA	-16.968	-10.052	5.756
1	MET HN1	-18.363	-11.296	4.808
1	MET HN2	-18.964	-10.642	6.192
1	MET HN3	-18.836	-9.728	4.811
1	MET HA	-17.022	-9.109	6.345
1	MET C	-16.058	-9.653	4.544
1	MET O	-15.491	-8.562	4.547
1	MET CB	-16.367	-11.107	6.732
1	MET HB1	-15.418	-10.692	7.133
1	MET HB2	-16.999	-11.196	7.638
1	MET CG	-16.048	-12.534	6.208
1	MET SD	-17.501	-13.438	5.624
1	MET CE	-18.199	-13.998	7.185
1	MET HG1	-15.303	-12.501	5.387
1	MET HG2	-15.555	-13.137	6.997
1	MET HE1	-19.124	-14.578	7.010
1	MET HE2	-17.490	-14.662	7.714
1	MET HE3	-18.445	-13.157	7.857
2	GLU N	-15.964	-10.507	3.518
2	GLU CA	-15.197	-10.254	2.269
2	GLU HN	-16.431	-11.415	3.613
2	GLU HA	-14.384	-9.526	2.462
2	GLU C	-16.086	-9.579	1.149
2	GLU O	-16.809	-8.614	1.422
2	GLU CB	-14.496	-11.624	1.943
2	GLU CG	-15.304	-12.969	1.947
2	GLU CD	-16.679	-12.989	1.312
2	GLU OE1	-16.770	-13.228	0.092

2	GLU	OE2	-17.689	-12.746	2.009
2	GLU	HB1	-13.980	-11.532	0.967
2	GLU	HB2	-13.654	-11.750	2.652
2	GLU	HG1	-14.713	-13.761	1.451
2	GLU	HG2	-15.430	-13.335	2.980
3	LYS	N	-16.030	-10.063	-0.108
3	LYS	CA	-16.999	-9.726	-1.196
3	LYS	HN	-15.594	-10.995	-0.096
3	LYS	HA	-16.597	-10.266	-2.076
3	LYS	C	-16.983	-8.231	-1.670
3	LYS	O	-16.139	-7.854	-2.488
3	LYS	CB	-18.379	-10.400	-0.889
3	LYS	CG	-19.212	-10.831	-2.119
3	LYS	CD	-20.381	-11.791	-1.776
3	LYS	CE	-20.044	-13.300	-1.688
3	LYS	NZ	-19.369	-13.685	-0.411
3	LYS	HB1	-18.203	-11.293	-0.268
3	LYS	HB2	-18.978	-9.753	-0.222
3	LYS	HG1	-19.611	-9.917	-2.608
3	LYS	HG2	-18.563	-11.290	-2.893
3	LYS	HD1	-20.925	-11.444	-0.874
3	LYS	HD2	-21.132	-11.674	-2.584
3	LYS	HE1	-20.986	-13.883	-1.807
3	LYS	HE2	-19.429	-13.605	-2.567
3	LYS	HZ1	-18.360	-13.394	-0.340
3	LYS	HZ2	-19.747	-13.259	0.445
3	LYS	HZ3	-19.300	-14.692	-0.225
4	VAL	N	-17.850	-7.370	-1.113
4	VAL	CA	-17.729	-5.883	-1.250
4	VAL	HN	-18.368	-7.786	-0.332
4	VAL	HA	-17.617	-5.678	-2.335
4	VAL	C	-16.444	-5.249	-0.593
4	VAL	O	-15.841	-4.361	-1.194
4	VAL	CB	-19.085	-5.211	-0.839

4	VAL	HB	-19.895	-5.756	-1.368
4	VAL	CG1	-19.192	-3.746	-1.314
4	VAL	HG11	-20.191	-3.313	-1.113
4	VAL	HG12	-18.450	-3.089	-0.822
4	VAL	HG13	-19.028	-3.652	-2.405
4	VAL	CG2	-19.423	-5.252	0.671
4	VAL	HG21	-20.429	-4.840	0.877
4	VAL	HG22	-19.412	-6.281	1.073
4	VAL	HG23	-18.707	-4.660	1.273
5	GLN	N	-15.991	-5.716	0.586
5	GLN	CA	-14.681	-5.298	1.177
5	GLN	HN	-16.525	-6.517	0.946
5	GLN	HA	-14.601	-4.196	1.093
5	GLN	C	-13.383	-5.831	0.471
5	GLN	O	-12.357	-5.141	0.491
5	GLN	CB	-14.696	-5.643	2.692
5	GLN	HB1	-13.684	-5.487	3.116
5	GLN	HB2	-14.878	-6.728	2.817
5	GLN	CG	-15.703	-4.852	3.575
5	GLN	HG1	-16.738	-4.985	3.209
5	GLN	HG2	-15.711	-5.300	4.586
5	GLN	CD	-15.396	-3.363	3.762
5	GLN	OE1	-14.647	-2.963	4.642
5	GLN	NE2	-15.948	-2.495	2.953
5	GLN	HE21	-15.630	-1.536	3.116
5	GLN	HE22	-16.412	-2.871	2.123
6	TYR	N	-13.412	-7.004	-0.197
6	TYR	CA	-12.392	-7.346	-1.233
6	TYR	HN	-14.349	-7.420	-0.207
6	TYR	HA	-11.396	-7.186	-0.771
6	TYR	C	-12.416	-6.413	-2.498
6	TYR	O	-11.364	-5.894	-2.875
6	TYR	CB	-12.491	-8.863	-1.566
6	TYR	HB1	-12.592	-9.455	-0.636

6	TYR	HB2	-13.426	-9.070	-2.123
6	TYR	CG	-11.270	-9.381	-2.345
6	TYR	CD1	-10.084	-9.668	-1.661
6	TYR	HD1	-10.045	-9.609	-0.583
6	TYR	CD2	-11.292	-9.461	-3.742
6	TYR	HD2	-12.200	-9.241	-4.287
6	TYR	CE1	-8.932	-10.001	-2.366
6	TYR	HE1	-8.027	-10.232	-1.830
6	TYR	CE2	-10.131	-9.785	-4.444
6	TYR	HE2	-10.139	-9.832	-5.524
6	TYR	CZ	-8.947	-10.036	-3.755
6	TYR	OH	-7.791	-10.280	-4.452
6	TYR	HH	-7.019	-10.219	-3.861
7	LEU	N	-13.587	-6.154	-3.113
7	LEU	CA	-13.759	-5.074	-4.135
7	LEU	HN	-14.394	-6.657	-2.722
7	LEU	HA	-13.139	-5.360	-5.005
7	LEU	C	-13.271	-3.631	-3.733
7	LEU	O	-12.609	-2.977	-4.542
7	LEU	CB	-15.245	-5.133	-4.594
7	LEU	HB1	-15.523	-6.180	-4.824
7	LEU	HB2	-15.880	-4.860	-3.731
7	LEU	CG	-15.651	-4.252	-5.804
7	LEU	HG	-15.360	-3.200	-5.603
7	LEU	CD1	-17.176	-4.286	-5.987
7	LEU	HD11	-17.553	-5.304	-6.199
7	LEU	HD12	-17.706	-3.921	-5.088
7	LEU	HD13	-17.502	-3.639	-6.825
7	LEU	CD2	-14.975	-4.703	-7.111
7	LEU	HD21	-15.191	-5.760	-7.352
7	LEU	HD22	-15.308	-4.098	-7.975
7	LEU	HD23	-13.876	-4.592	-7.065
8	THR	N	-13.511	-3.164	-2.491
8	THR	CA	-12.882	-1.929	-1.922

8	THR	HN	-14.206	-3.711	-1.961
8	THR	HA	-13.236	-1.074	-2.530
8	THR	C	-11.312	-1.892	-2.013
8	THR	O	-10.767	-0.964	-2.613
8	THR	CB	-13.426	-1.715	-0.469
8	THR	OG1	-14.849	-1.671	-0.441
8	THR	HB	-13.095	-2.565	0.160
8	THR	HG1	-15.101	-0.841	-0.857
8	THR	CG2	-12.988	-0.414	0.216
8	THR	HG21	-13.230	0.484	-0.384
8	THR	HG22	-13.473	-0.292	1.203
8	THR	HG23	-11.898	-0.399	0.403
9	ARG	N	-10.582	-2.897	-1.486
9	ARG	CA	-9.100	-2.999	-1.688
9	ARG	HN	-11.154	-3.657	-1.099
9	ARG	HA	-8.665	-2.002	-1.470
9	ARG	C	-8.597	-3.341	-3.141
9	ARG	O	-7.489	-2.943	-3.504
9	ARG	CB	-8.480	-3.985	-0.662
9	ARG	CG	-8.679	-3.663	0.845
9	ARG	CD	-7.677	-4.331	1.815
9	ARG	NE	-7.723	-5.828	1.787
9	ARG	CZ	-6.925	-6.617	1.073
9	ARG	NH1	-7.116	-7.891	1.115
9	ARG	NH2	-5.961	-6.190	0.314
9	ARG	HB1	-7.392	-4.020	-0.865
9	ARG	HB2	-8.843	-5.012	-0.870
9	ARG	HG1	-8.611	-2.565	0.992
9	ARG	HG2	-9.719	-3.911	1.140
9	ARG	HD1	-6.654	-3.934	1.650
9	ARG	HD2	-7.914	-3.993	2.846
9	ARG	HE	-8.426	-6.349	2.319
9	ARG	HH12	-7.864	-8.264	1.693
9	ARG	HH11	-6.571	-8.401	0.386

9	ARG	HH21	-5.791	-5.188	0.334
9	ARG	HH22	-5.373	-6.910	-0.147
10	SER	N	-9.350	-4.086	-3.965
10	SER	CA	-9.062	-4.255	-5.423
10	SER	HN	-10.216	-4.451	-3.541
10	SER	HA	-7.998	-4.514	-5.543
10	SER	C	-9.237	-2.985	-6.326
10	SER	O	-8.384	-2.744	-7.184
10	SER	CB	-9.828	-5.503	-5.913
10	SER	OG	-9.313	-6.678	-5.275
10	SER	HB1	-10.914	-5.399	-5.724
10	SER	HB2	-9.719	-5.609	-7.011
10	SER	HG	-9.884	-7.422	-5.502
11	ALA	N	-10.260	-2.142	-6.107
11	ALA	CA	-10.294	-0.755	-6.648
11	ALA	HN	-10.941	-2.461	-5.403
11	ALA	HA	-10.105	-0.797	-7.740
11	ALA	C	-9.225	0.239	-6.059
11	ALA	O	-8.524	0.906	-6.827
11	ALA	CB	-11.743	-0.266	-6.454
11	ALA	HB1	-12.024	-0.198	-5.385
11	ALA	HB2	-11.891	0.739	-6.892
11	ALA	HB3	-12.479	-0.934	-6.942
12	ILE	N	-9.063	0.325	-4.722
12	ILE	CA	-8.073	1.248	-4.069
12	ILE	HN	-9.733	-0.232	-4.175
12	ILE	HA	-8.213	2.243	-4.537
12	ILE	C	-6.563	0.911	-4.365
12	ILE	O	-5.785	1.854	-4.502
12	ILE	CB	-8.421	1.468	-2.546
12	ILE	HB	-8.473	0.468	-2.073
12	ILE	CG1	-9.804	2.172	-2.347
12	ILE	HG11	-10.558	1.710	-3.012
12	ILE	HG12	-9.748	3.226	-2.688

12 ILE	CG2	-7.349	2.290	-1.770
12 ILE	HG21	-7.590	2.410	-0.699
12 ILE	HG22	-6.350	1.816	-1.796
12 ILE	HG23	-7.222	3.309	-2.187
12 ILE	CD1	-10.379	2.121	-0.921
12 ILE	HD11	-11.413	2.512	-0.890
12 ILE	HD12	-10.405	1.088	-0.531
12 ILE	HD13	-9.790	2.730	-0.211
13 ARG	N	-6.116	-0.357	-4.492
13 ARG	CA	-4.688	-0.677	-4.826
13 ARG	HN	-6.837	-1.075	-4.352
13 ARG	HA	-4.087	-0.227	-4.011
13 ARG	C	-4.064	-0.040	-6.124
13 ARG	O	-2.892	0.342	-6.094
13 ARG	CB	-4.446	-2.208	-4.706
13 ARG	CG	-4.985	-3.125	-5.840
13 ARG	CD	-4.694	-4.625	-5.629
13 ARG	NE	-5.635	-5.210	-4.631
13 ARG	CZ	-5.642	-6.473	-4.222
13 ARG	NH1	-4.693	-7.326	-4.456
13 ARG	NH2	-6.651	-6.890	-3.537
13 ARG	HB1	-4.825	-2.556	-3.726
13 ARG	HB2	-3.351	-2.360	-4.640
13 ARG	HG1	-6.065	-2.948	-6.011
13 ARG	HG2	-4.505	-2.820	-6.792
13 ARG	HD1	-4.788	-5.161	-6.598
13 ARG	HD2	-3.634	-4.753	-5.333
13 ARG	HE	-6.427	-4.657	-4.281
13 ARG	HH12	-3.840	-6.971	-4.905
13 ARG	HH11	-4.827	-8.262	-4.029
13 ARG	HH21	-7.482	-6.302	-3.505
13 ARG	HH22	-6.560	-7.869	-3.213
14 ARG	N	-4.825	0.102	-7.229
14 ARG	CA	-4.376	0.917	-8.401

14 ARG	HN	-5.808	-0.154	-7.074
14 ARG	HA	-3.272	0.844	-8.428
14 ARG	C	-4.656	2.464	-8.334
14 ARG	O	-3.866	3.230	-8.890
14 ARG	CB	-4.845	0.244	-9.719
14 ARG	CG	-4.019	0.701	-10.958
14 ARG	CD	-4.063	-0.202	-12.216
14 ARG	NE	-3.844	-1.670	-12.010
14 ARG	CZ	-2.795	-2.247	-11.427
14 ARG	NH1	-1.732	-1.615	-11.038
14 ARG	NH2	-2.842	-3.520	-11.225
14 ARG	HB1	-4.739	-0.853	-9.604
14 ARG	HB2	-5.927	0.412	-9.882
14 ARG	HG1	-4.336	1.727	-11.239
14 ARG	HG2	-2.961	0.842	-10.667
14 ARG	HD1	-5.052	-0.063	-12.698
14 ARG	HD2	-3.349	0.185	-12.973
14 ARG	HE	-4.569	-2.354	-12.237
14 ARG	HH12	-1.739	-0.612	-11.189
14 ARG	HH11	-1.056	-2.146	-10.443
14 ARG	HH21	-3.658	-4.047	-11.522
14 ARG	HH22	-2.019	-3.907	-10.697
15 ALA	N	-5.709	2.944	-7.642
15 ALA	CA	-5.801	4.374	-7.222
15 ALA	HN	-6.269	2.220	-7.179
15 ALA	HA	-5.765	5.005	-8.134
15 ALA	C	-4.644	4.893	-6.290
15 ALA	O	-4.074	5.946	-6.571
15 ALA	CB	-7.199	4.561	-6.603
15 ALA	HB1	-7.394	5.627	-6.376
15 ALA	HB2	-8.010	4.227	-7.279
15 ALA	HB3	-7.305	4.011	-5.651
16 SER	N	-4.250	4.134	-5.248
16 SER	CA	-3.106	4.482	-4.352

16 SER HN	-4.848	3.315	-5.069
16 SER HA	-3.235	5.535	-4.049
16 SER C	-1.638	4.397	-4.935
16 SER O	-0.702	4.880	-4.297
16 SER CB	-3.324	3.676	-3.050
16 SER OG	-2.393	4.057	-2.031
16 SER HB1	-3.255	2.589	-3.240
16 SER HB2	-4.345	3.849	-2.656
16 SER HG	-2.450	5.037	-1.916
17 THR N	-1.403	3.876	-6.157
17 THR CA	-0.201	4.263	-6.976
17 THR HN	-2.258	3.512	-6.589
17 THR HA	0.636	4.460	-6.276
17 THR C	-0.319	5.601	-7.802
17 THR O	0.712	6.218	-8.084
17 THR CB	0.310	3.069	-7.842
17 THR OG1	1.531	3.425	-8.481
17 THR HB	0.520	2.218	-7.163
17 THR HG1	1.382	4.296	-8.872
17 THR CG2	-0.615	2.566	-8.957
17 THR HG21	-0.136	1.754	-9.534
17 THR HG22	-1.545	2.157	-8.534
17 THR HG23	-0.896	3.364	-9.669
18 ILE N	-1.523	6.028	-8.226
18 ILE CA	-1.764	7.358	-8.891
18 ILE HN	-2.302	5.493	-7.824
18 ILE HA	-0.928	7.554	-9.592
18 ILE C	-1.733	8.562	-7.864
18 ILE O	-1.063	9.563	-8.116
18 ILE CB	-3.077	7.284	-9.765
18 ILE HB	-3.906	7.053	-9.065
18 ILE CG1	-3.060	6.171	-10.865
18 ILE HG11	-2.576	5.258	-10.472
18 ILE HG12	-2.417	6.477	-11.714

18 ILE	CG2	-3.416	8.637	-10.450
18 ILE	HG21	-3.540	9.455	-9.715
18 ILE	HG22	-2.626	8.959	-11.154
18 ILE	HG23	-4.364	8.600	-11.016
18 ILE	CD1	-4.448	5.738	-11.380
18 ILE	HD11	-4.987	6.560	-11.884
18 ILE	HD12	-4.365	4.911	-12.110
18 ILE	HD13	-5.090	5.375	-10.555
19 GLU	N	-2.426	8.437	-6.718
19 GLU	CA	-2.209	9.220	-5.456
19 GLU	HN	-2.954	7.556	-6.669
19 GLU	HA	-2.898	10.084	-5.466
19 GLU	C	-0.786	9.777	-5.056
19 GLU	O	-0.692	10.794	-4.362
19 GLU	CB °	-2.714	8.196	-4.390
19 GLU	CG	-2.806	8.605	-2.896
19 GLU	CD	-3.194	7.442	-1.992
19 GLU	OE1	-2.326	6.599	-1.686
19 GLU	OE2	-4.375	7.339	-1.613
19 GLU	HB1	-3.720	7.818	-4.667
19 GLU	HB2	-2.044	7.314	-4.436
19 GLU	HG1	-1.845	8.993	-2.521
19 GLU	HG2	-3.546	9.409	-2.747
20 MET	N	0.304	9.081	-5.411
20 MET	CA	1.613	9.205	-4.724
20 MET	HN	0.064	8.269	-5.989
20 MET	HA	1.499	9.758	-3.773
20 MET	C	2.703	9.961	-5.580
20 MET	O	3.391	9.307	-6.372
20 MET	CB	2.021	7.758	-4.320
20 MET	HB1	1.924	7.072	-5.187
20 MET	HB2	3.096	7.734	-4.080
20 MET	CG	1.245	7.190	-3.111
20 MET	SD	1.761	5.508	-2.777

20 MET CE	0.862	5.232	-1.245
20 MET HG1	1.366	7.818	-2.209
20 MET HG2	0.155	7.174	-3.311
20 MET HE1	0.982	4.201	-0.876
20 MET HE2	1.153	5.949	-0.457
20 MET HE3	-0.219	5.377	-1.430
21 PRO N	2.965	11.301	-5.432
21 PRO CA	4.032	12.006	-6.211
21 PRO CD	2.094	12.217	-4.662
21 PRO HA	3.958	11.712	-7.278
21 PRO HD1	1.931	11.903	-3.614
21 PRO HD2	1.100	12.301	-5.145
21 PRO C	5.513	11.694	-5.772
21 PRO O	5.777	10.728	-5.055
21 PRO CB	3.558	13.475	-6.078
21 PRO HB1	4.365	14.226	-6.161
21 PRO HB2	2.843	13.707	-6.894
21 PRO CG	2.833	13.550	-4.735
21 PRO HG1	2.144	14.412	-4.664
21 PRO HG2	3.555	13.637	-3.900
22 GLN N	6.492	12.482	-6.259
22 GLN CA	7.971	12.242	-6.117
22 GLN HN	6.130	13.269	-6.807
22 GLN HA	8.237	11.652	-7.014
22 GLN C	8.571	11.380	-4.938
22 GLN O	9.183	10.336	-5.185
22 GLN CB	8.687	13.611	-6.328
22 GLN HB1	9.776	13.432	-6.440
22 GLN HB2	8.393	14.021	-7.315
22 GLN CG	8.480	14.719	-5.249
22 GLN HG1	7.400	14.880	-5.075
22 GLN HG2	8.893	14.382	-4.280
22 GLN CD	9.089	16.086	-5.575
22 GLN OE1	9.447	16.412	-6.700

22	GLN	NE2	9.203	16.949	-4.600
22	GLN	HE21	8.880	16.664	-3.672
22	GLN	HE22	9.586	17.853	-4.892
23	GLN	N	8.363	11.795	-3.682
23	GLN	CA	8.769	11.029	-2.462
23	GLN	HN	7.850	12.681	-3.626
23	GLN	HA	9.766	10.583	-2.655
23	GLN	C	7.858	9.817	-2.049
23	GLN	O	8.366	8.773	-1.628
23	GLN	CB	8.984	12.075	-1.331
23	GLN	HB1	9.753	12.795	-1.680
23	GLN	HB2	9.473	11.581	-0.469
23	GLN	CG	7.730	12.848	-0.815
23	GLN	HG1	6.971	12.936	-1.613
23	GLN	HG2	7.219	12.273	-0.019
23	GLN	CD	8.018	14.281	-0.367
23	GLN	OE1	8.162	15.184	-1.181
23	GLN	NE2	8.113	14.540	0.911
23	GLN	HE21	7.995	13.761	1.562
23	GLN	HE22	8.301	15.524	1.123
24	ALA	N	6.527	9.922	-2.191
24	ALA	CA	5.598	8.770	-2.032
24	ALA	HN	6.246	10.756	-2.719
24	ALA	HA	5.810	8.295	-1.054
24	ALA	C	5.702	7.618	-3.104
24	ALA	O	5.540	6.445	-2.745
24	ALA	CB	4.202	9.401	-1.926
24	ALA	HB1	4.133	10.148	-1.112
24	ALA	HB2	3.901	9.909	-2.861
24	ALA	HB3	3.433	8.639	-1.708
25	ARG	N	6.020	7.917	-4.378
25	ARG	CA	6.491	6.901	-5.368
25	ARG	HN	5.937	8.921	-4.609
25	ARG	HA	5.870	6.003	-5.211

25 ARG C	7.958	6.348	-5.212
25 ARG O	8.226	5.246	-5.698
25 ARG CB	6.139	7.358	-6.810
25 ARG CG	6.910	8.577	-7.389
25 ARG CD	7.630	8.344	-8.738
25 ARG NE	8.675	7.276	-8.682
25 ARG CZ	9.882	7.391	-8.144
25 ARG NH1	10.343	8.483	-7.617
25 ARG NH2	10.647	6.350	-8.127
25 ARG HB1	5.050	7.561	-6.864
25 ARG HB2	6.250	6.485	-7.481
25 ARG HG1	6.185	9.408	-7.513
25 ARG HG2	7.623	8.968	-6.640
25 ARG HD1	6.870	8.070	-9.502
25 ARG HD2	8.049	9.297	-9.123
25 ARG HE	8.487	6.333	-9.036
25 ARG HH12	9.678	9.253	-7.562
25 ARG HH11	11.269	8.411	-7.176
25 ARG HH21	10.262	5.474	-8.478
25 ARG HH22	11.531	6.451	-7.608
26 GLN N	8.883	7.013	-4.478
26 GLN CA	10.024	6.300	-3.813
26 GLN HN	8.547	7.923	-4.141
26 GLN HA	10.567	5.729	-4.593
26 GLN C	9.557	5.221	-2.773
26 GLN O	9.940	4.063	-2.908
26 GLN CB	11.056	7.278	-3.178
26 GLN HB1	10.541	7.954	-2.472
26 GLN HB2	11.746	6.698	-2.531
26 GLN CG	11.938	8.128	-4.132
26 GLN HG1	12.574	8.807	-3.533
26 GLN HG2	11.290	8.808	-4.715
26 GLN CD	12.851	7.379	-5.112
26 GLN OE1	12.666	7.434	-6.322

26 GLN	NE2	13.859	6.683	-4.658
26 GLN	HE21	13.951	6.566	-3.644
26 GLN	HE22	14.393	6.190	-5.379
27 LYS	N	8.661	5.530	-1.814
27 LYS	CA	7.965	4.488	-0.985
27 LYS	HN	8.437	6.532	-1.767
27 LYS	HA	8.749	3.968	-0.397
27 LYS	C	7.205	3.320	-1.738
27 LYS	O	6.925	2.294	-1.118
27 LYS	CB	7.084	5.278	0.032
27 LYS	CG	6.375	4.464	1.151
27 LYS	CD	4.824	4.447	1.123
27 LYS	CE	4.146	3.508	0.100
27 LYS	NZ	4.039	4.152	-1.240
27 LYS	HB1	6.354	5.914	-0.500
27 LYS	HB2	7.736	6.020	0.539
27 LYS	HG1	6.775	3.431	1.209
27 LYS	HG2	6.683	4.897	2.125
27 LYS	HD1	4.499	4.110	2.129
27 LYS	HD2	4.421	5.479	1.064
27 LYS	HE1	4.692	2.536	0.045
27 LYS	HE2	3.132	3.226	0.470
27 LYS	HZ1	3.433	4.985	-1.252
27 LYS	HZ2	4.943	4.472	-1.617
27 LYS	HZ3	3.653	3.537	-1.972
28 LEU	N	6.852	3.447	-3.034
28 LEU	CA	6.417	2.288	-3.875
28 LEU	HN	7.273	4.279	-3.462
28 LEU	HA	5.700	1.669	-3.298
28 LEU	C	7.599	1.309	-4.230
28 LEU	O	7.644	0.200	-3.695
28 LEU	CB	5.656	2.809	-5.135
28 LEU	HB1	6.358	3.414	-5.737
28 LEU	HB2	5.429	1.941	-5.788

28 LEU	CG	4.345	3.618	-4.944
28 LEU	HG	4.521	4.455	-4.242
28 LEU	CD1	3.889	4.207	-6.289
28 LEU	HD11	3.626	3.417	-7.017
28 LEU	HD12	2.996	4.851	-6.178
28 LEU	HD13	4.664	4.825	-6.770
28 LEU	CD2	3.199	2.752	-4.415
28 LEU	HD21	2.261	3.324	-4.268
28 LEU	HD22	2.945	1.949	-5.132
28 LEU	HD23	3.435	2.251	-3.461
29 GLN	N	8.551	1.701	-5.102
29 GLN	CA	9.681	0.813	-5.525
29 GLN	HN	8.445	2.659	-5.444
29 GLN	HA	9.296	-0.228	-5.547
29 GLN	C	10.927	0.704	-4.581
29 GLN	O	11.555	-0.350	-4.563
29 GLN	CB	10.058	1.100	-7.008
29 GLN	HB1	10.824	0.361	-7.320
29 GLN	HB2	9.184	0.826	-7.631
29 GLN	CG	10.521	2.525	-7.440
29 GLN	HG1	9.707	3.243	-7.230
29 GLN	HG2	10.628	2.539	-8.541
29 GLN	CD	11.813	3.109	-6.855
29 GLN	OE1	11.817	4.184	-6.267
29 GLN	NE2	12.940	2.471	-7.031
29 GLN	HE21	12.865	1.517	-7.397
29 GLN	HE22	13.731	2.870	-6.518
30 ASN	N	11.300	1.729	-3.796
30 ASN	CA	12.455	1.663	-2.841
30 ASN	HN	10.673	2.543	-3.835
30 ASN	HA	13.328	1.287	-3.409
30 ASN	C	12.294	0.683	-1.616
30 ASN	O	13.296	0.164	-1.122
30 ASN	CB	12.777	3.128	-2.412

30 ASN HB1	12.124	3.435	-1.575
30 ASN HB2	12.518	3.827	-3.229
30 ASN CG	14.217	3.499	-2.051
30 ASN OD1	14.740	4.505	-2.512
30 ASN ND2	14.897	2.774	-1.201
30 ASN HD21	14.495	1.861	-0.955
30 ASN HD22	15.872	3.070	-1.104
31 LEU N	11.065	0.414	-1.141
31 LEU CA	10.771	-0.779	-0.283
31 LEU HN	10.324	0.878	-1.678
31 LEU HA	11.547	-0.852	0.505
31 LEU C	10.791	-2.187	-0.999
31 LEU O	11.021	-3.200	-0.334
31 LEU CB	9.400	-0.555	0.422
31 LEU HB1	8.613	-0.511	-0.357
31 LEU HB2	9.161	-1.468	1.000
31 LEU CG	9.234	0.657	1.379
31 LEU HG	9.418	1.589	0.808
31 LEU CD1	7.793	0.696	1.913
31 LEU HD11	7.540	-0.215	2.491
31 LEU HD12	7.624	1.560	2.582
31 LEU HD13	7.056	0.771	1.091
31 LEU CD2	10.205	0.614	2.572
31 LEU HD21	10.050	1.463	3.263
31 LEU HD22	10.099	-0.315	3.165
31 LEU HD23	11.260	0.669	2.244
32 PHE N	10.538	-2.258	-2.316
32 PHE CA	10.696	-3.495	-3.136
32 PHE HN	10.482	-1.340	-2.768
32 PHE HA	10.395	-4.364	-2.515
32 PHE C	12.179	-3.811	-3.566
32 PHE O	12.658	-4.914	-3.301
32 PHE CB	9.693	-3.392	-4.333
32 PHE HB1	9.145	-2.429	-4.348

32 PHE	HB2	10.258	-3.342	-5.286
32 PHE	CG	8.648	-4.516	-4.425
32 PHE	CD1	8.796	-5.536	-5.370
32 PHE	HD1	9.665	-5.563	-6.012
32 PHE	CD2	7.508	-4.495	-3.613
32 PHE	HD2	7.370	-3.707	-2.887
32 PHE	CE1	7.815	-6.513	-5.511
32 PHE	HE1	7.935	-7.287	-6.256
32 PHE	CE2	6.530	-5.480	-3.751
32 PHE	HE2	5.644	-5.463	-3.137
32 PHE	CZ	6.679	-6.481	-4.707
32 PHE	HZ	5.903	-7.220	-4.840
33 ILE	N	12.931	-2.860	-4.163
33 ILE	CA	14.249	-3.145	-4.844
33 ILE	HN	12.381	-2.034	-4.432
33 ILE	HA	14.130	-4.126	-5.346
33 ILE	C	15.529	-3.369	-3.934
33 ILE	O	16.676	-3.233	-4.369
33 ILE	CB	14.418	-2.106	-6.019
33 ILE	HB	13.426	-2.035	-6.516
33 ILE	CG1	15.384	-2.601	-7.135
33 ILE	HG11	15.188	-3.677	-7.326
33 ILE	HG12	16.434	-2.566	-6.789
33 ILE	CG2	14.822	-0.685	-5.541
33 ILE	HG21	14.801	0.051	-6.363
33 ILE	HG22	14.142	-0.317	-4.756
33 ILE	HG23	15.844	-0.676	-5.117
33 ILE	CD1	15.254	-1.881	-8.489
33 ILE	HD11	14.229	-1.989	-8.896
33 ILE	HD12	15.491	-0.806	-8.425
33 ILE	HD13	15.936	-2.326	-9.236
34 ASN	N	15.344	-3.823	-2.688
34 ASN	CA	16.279	-4.802	-2.047
34 ASN	HN	14.332	-3.912	-2.532

34 ASN	HA	17.316	-4.527	-2.325
34 ASN	C	16.140	-6.303	-2.523
34 ASN	O	17.115	-7.053	-2.449
34 ASN	CB	16.199	-4.624	-0.503
34 ASN	HB1	16.971	-5.266	-0.040
34 ASN	HB2	16.514	-3.598	-0.235
34 ASN	CG	14.875	-4.948	0.199
34 ASN	OD1	14.645	-6.042	0.695
34 ASN	ND2	13.967	-4.009	0.275
34 ASN	HD21	14.152	-3.184	-0.301
34 ASN	HD22	13.053	-4.355	0.578
35 PHE	N	14.955	-6.729	-2.996
35 PHE	CA	14.720	-8.055	-3.644
35 PHE	HN	14.230	-6.001	-3.022
35 PHE	HA	15.681	-8.604	-3.688
35 PHE	C	14.267	-7.997	-5.150
35 PHE	O	14.703	-8.857	-5.922
35 PHE	CB	13.790	-8.914	-2.732
35 PHE	HB1	14.297	-9.051	-1.757
35 PHE	HB2	13.757	-9.938	-3.149
35 PHE	CG	12.346	-8.419	-2.481
35 PHE	CD1	11.297	-8.823	-3.315
35 PHE	HD1	11.490	-9.483	-4.149
35 PHE	CD2	12.085	-7.518	-1.443
35 PHE	HD2	12.890	-7.171	-0.812
35 PHE	CE1	10.013	-8.316	-3.124
35 PHE	HE1	9.216	-8.591	-3.799
35 PHE	CE2	10.805	-7.002	-1.262
35 PHE	HE2	10.636	-6.252	-0.503
35 PHE	CZ	9.770	-7.402	-2.104
35 PHE	HZ	8.787	-6.965	-2.001
36 CYS	N	13.391	-7.059	-5.568
36 CYS	CA	12.937	-6.919	-6.979
36 CYS	C	12.642	-5.431	-7.337

36	CYS	O	13.077	-4.984	-8.423
36	CYS	CB	11.711	-7.838	-7.186
36	CYS	SG	11.181	-7.821	-8.935
36	CYS	OXT	11.882	-4.739	-6.619
36	CYS	HN	13.082	-6.376	-4.869
36	CYS	HA	13.741	-7.243	-7.669
36	CYS	HB1	11.954	-8.878	-6.906
36	CYS	HB2	10.872	-7.519	-6.542
36	CYS	HG	10.855	-6.530	-9.009
END					
1	CYS	N	8.859	-12.826	-0.866
1	CYS	CA	7.463	-12.830	-1.374
1	CYS	HN1	9.334	-13.726	-0.991
1	CYS	HN2	9.421	-12.087	-1.305
1	CYS	HN3	8.864	-12.628	0.142
1	CYS	HA	7.468	-13.235	-2.407
1	CYS	C	6.828	-11.396	-1.494
1	CYS	O	7.522	-10.392	-1.317
1	CYS	CB	6.712	-13.837	-0.472
1	CYS	SG	6.308	-13.188	1.176
1	CYS	HB1	5.744	-14.101	-0.939
1	CYS	HB2	7.235	-14.807	-0.360
2	TYR	N	5.529	-11.294	-1.812
2	TYR	CA	4.750	-10.029	-1.672
2	TYR	HN	5.009	-12.171	-1.934
2	TYR	HA	5.241	-9.395	-0.905
2	TYR	C	3.318	-10.348	-1.118
2	TYR	O	2.710	-11.362	-1.482
2	TYR	CB	4.792	-9.194	-2.987
2	TYR	HB1	4.452	-8.169	-2.757
2	TYR	HB2	5.851	-9.037	-3.272
2	TYR	CG	3.999	-9.693	-4.206
2	TYR	CD1	4.589	-10.559	-5.132
2	TYR	HD1	5.609	-10.885	-5.003

2	TYR	CD2	2.682	-9.266	-4.405
2	TYR	HD2	2.215	-8.590	-3.702
2	TYR	CE1	3.859	-11.011	-6.232
2	TYR	HE1	4.307	-11.690	-6.943
2	TYR	CE2	1.957	-9.716	-5.506
2	TYR	HE2	0.936	-9.388	-5.641
2	TYR	CZ	2.547	-10.590	-6.416
2	TYR	OH	1.833	-11.038	-7.494
2	TYR	HH	0.967	-10.629	-7.471
3	TRP	N	2.784	-9.498	-0.230
3	TRP	CA	1.456	-9.732	0.411
3	TRP	HN	3.400	-8.720	0.047
3	TRP	HA	1.393	-10.807	0.685
3	TRP	C	0.227	-9.448	-0.539
3	TRP	O	0.394	-9.065	-1.703
3	TRP	CB	1.498	-8.898	1.727
3	TRP	HB1	2.506	-8.943	2.193
3	TRP	HB2	1.353	-7.829	1.491
3	TRP	CG	0.519	-9.345	2.820
3	TRP	CD1	-0.639	-8.655	3.245
3	TRP	CD2	0.588	-10.478	3.609
3	TRP	NE1	-1.306	-9.343	4.282
3	TRP	CE2	-0.520	-10.467	4.493
3	TRP	HD1	-0.967	-7.716	2.824
3	TRP	HE1	-2.095	-9.033	4.857
3	TRP	CE3	1.538	-11.532	3.661
3	TRP	HE3	2.409	-11.541	3.020
3	TRP	CZ2	-0.682	-11.505	5.441
3	TRP	HZ2	-1.510	-11.494	6.133
3	TRP	CZ3	1.353	-12.549	4.598
3	TRP	HZ3	2.088	-13.342	4.654
3	TRP	CH2	0.260	-12.537	5.476
3	TRP	HH2	0.160	-13.332	6.200
4	GLU	N	-1.026	-9.624	-0.065

4	GLU	CA	-2.251	-9.398	-0.900
4	GLU	HN	-1.066	-9.985	0.896
4	GLU	HA	-2.059	-9.850	-1.895
4	GLU	C	-2.579	-7.874	-1.150
4	GLU	O	-3.531	-7.282	-0.629
4	GLU	CB	-3.423	-10.225	-0.289
4	GLU	CG	-4.548	-10.658	-1.279
4	GLU	CD	-5.428	-9.594	-1.902
4	GLU	OE1	-5.983	-8.751	-1.176
4	GLU	OE2	-5.630	-9.609	-3.130
4	GLU	HB1	-3.026	-11.161	0.155
4	GLU	HB2	-3.858	-9.678	0.569
4	GLU	HG1	-4.114	-11.254	-2.103
4	GLU	HG2	-5.235	-11.354	-0.770
5	LEU	N	-1.753	-7.242	-1.991
5	LEU	CA	-1.972	-5.882	-2.541
5	LEU	HN	-0.922	-7.815	-2.212
5	LEU	HA	-3.057	-5.734	-2.704
5	LEU	C	-1.333	-5.919	-3.973
5	LEU	O	-1.967	-6.436	-4.899
5	LEU	CB	-1.517	-4.805	-1.497
5	LEU	HB1	-0.421	-4.840	-1.373
5	LEU	HB2	-1.897	-5.131	-0.508
5	LEU	CG	-2.002	-3.346	-1.723
5	LEU	HG	-3.062	-3.370	-2.045
5	LEU	CD1	-1.930	-2.530	-0.421
5	LEU	HD11	-2.500	-2.998	0.402
5	LEU	HD12	-0.889	-2.404	-0.065
5	LEU	HD13	-2.344	-1.512	-0.553
5	LEU	CD2	-1.183	-2.584	-2.778
5	LEU	HD21	-1.272	-3.021	-3.788
5	LEU	HD22	-1.512	-1.532	-2.878
5	LEU	HD23	-0.102	-2.555	-2.533
6	GLU	N	-0.102	-5.418	-4.147

6	GLU	CA	0.670	-5.433	-5.427
6	GLU	HN	0.318	-5.068	-3.279
6	GLU	HA	0.760	-6.480	-5.776
6	GLU	C	2.125	-4.958	-5.064
6	GLU	O	3.021	-5.761	-4.797
6	GLU	CB	-0.062	-4.615	-6.547
6	GLU	CG	0.643	-4.580	-7.927
6	GLU	CD	-0.149	-3.863	-8.997
6	GLU	OE1	-0.877	-4.532	-9.753
6	GLU	OE2	-0.072	-2.625	-9.100
6	GLU	HB1	-1.073	-5.034	-6.699
6	GLU	HB2	-0.241	-3.577	-6.203
6	GLU	HG1	1.630	-4.089	-7.869
6	GLU	HG2	0.842	-5.603	-8.299
8	TRP	N	2.313	-3.638	-4.926
8	TRP	CA	3.537	-2.986	-4.364
8	TRP	HN	1.515	-3.090	-5.263
8	TRP	HA	4.424	-3.433	-4.856
8	TRP	C	3.848	-3.117	-2.821
8	TRP	O	4.653	-2.346	-2.288
8	TRP	CB	3.456	-1.496	-4.839
8	TRP	HB1	4.309	-0.929	-4.424
8	TRP	HB2	3.659	-1.485	-5.927
8	TRP	CG	2.153	-0.690	-4.582
8	TRP	CD1	1.178	-0.428	-5.575
8	TRP	CD2	1.662	-0.105	-3.421
8	TRP	NE1	0.101	0.325	-5.068
8	TRP	CE2	0.430	0.525	-3.742
8	TRP	HD1	1.240	-0.780	-6.598
8	TRP	HE1	-0.765	0.609	-5.546
8	TRP	CE3	2.173	-0.038	-2.098
8	TRP	HE3	3.108	-0.507	-1.834
8	TRP	CZ2	-0.283	1.234	-2.751
8	TRP	HZ2	-1.204	1.736	-2.998

8	TRP	CZ3	1.437	0.648	-1.130
8	TRP	HZ3	1.792	0.681	-0.112
8	TRP	CH2	0.230	1.282	-1.455
8	TRP	HH2	-0.324	1.816	-0.694
8	LEU	N	3.264	-4.088	-2.102
8	LEU	CA	3.536	-4.329	-0.654
8	LEU	HN	2.865	-4.819	-2.705
8	LEU	HA	3.941	-3.411	-0.189
8	LEU	C	4.558	-5.516	-0.479
8	LEU	O	4.209	-6.629	-0.884
8	LEU	CB	2.165	-4.593	0.042
8	LEU	HB1	1.408	-3.926	-0.417
8	LEU	HB2	1.824	-5.619	-0.207
8	LEU	CG	2.061	-4.358	1.574
8	LEU	HG	2.296	-3.294	1.781
8	LEU	CD1	3.002	-5.228	2.425
8	LEU	HD11	4.062	-4.953	2.285
8	LEU	HD12	2.805	-5.135	3.510
8	LEU	HD13	2.927	-6.303	2.177
8	LEU	CD2	0.619	-4.623	2.040
8	LEU	HD21	0.492	-4.415	3.118
8	LEU	HD22	-0.111	-3.989	1.507
8	LEU	HD23	0.316	-5.674	1.876
9	PRO	N	5.780	-5.358	0.119
9	PRO	CA	6.694	-6.502	0.430
9	PRO	CD	6.411	-4.035	0.314
9	PRO	HA	7.030	-6.917	-0.541
9	PRO	HD1	6.017	-3.549	1.227
9	PRO	HD2	6.254	-3.346	-0.539
9	PRO	C	6.165	-7.738	1.254
9	PRO	O	4.962	-7.953	1.399
9	PRO	CB	7.873	-5.753	1.096
9	PRO	HB1	8.834	-6.283	0.969
9	PRO	HB2	7.706	-5.659	2.189

9	PRO	CG	7.892	-4.373	0.455
9	PRO	HG1	8.381	-4.416	-0.539
9	PRO	HG2	8.450	-3.636	1.059
10	CYS	N	7.083	-8.610	1.703
10	CYS	CA	6.766	-10.017	2.059
10	CYS	HN	8.045	-8.317	1.519
10	CYS	HA	6.123	-10.428	1.255
10	CYS	C	5.908	-10.281	3.352
10	CYS	O	4.684	-10.146	3.295
10	CYS	CB	8.136	-10.743	1.930
10	CYS	SG	7.929	-12.525	2.158
10	CYS	HB1	8.849	-10.378	2.694
10	CYS	HB2	8.613	-10.553	0.953
11	ALA	N	6.523	-10.747	4.456
11	ALA	CA	5.941	-11.825	5.306
11	ALA	HN	7.531	-10.788	4.308
11	ALA	HA	6.701	-12.037	6.082
11	ALA	C	5.832	-13.184	4.541
11	ALA	O	6.858	-13.901	4.464
11	ALA	OXT	4.767	-13.516	3.972
11	ALA	CB	4.694	-11.381	6.092
11	ALA	HB1	3.843	-11.199	5.412
11	ALA	HB2	4.376	-12.169	6.797
11	ALA	HB3	4.873	-10.456	6.664

END

Table V

Residue	NH	C _α H	C _β H	C _γ H	C _δ H	Others
Met1		4.159	2.124	2.566		2.012
Glu2	8.803	4.572	1.967	2.510, 2.319		
Lys3	8.787	4.147	1.900	1.559, 1.431	1.729	2.969
Val4	8.314	3.889	2.132	1.026, 0.951		
Gln5	7.846	4.092	2.364, <u>2.252</u>	2.369		7.431, 6.823
Tyr6	7.924	4.225	3.114		7.046	6.780
Leu7	8.536	4.013	1.831	1.586	0.915	
Thr8	7.975	3.955	4.247	1.222		
Arg9	8.156	3.518	1.845, 1.725	1.587	3.092	7.242
Ser10	7.991	4.082	<u>3.936</u> , 3.767			
Ala11	8.051	4.085	1.510			
Ile12	8.126	3.681	<u>1.857</u>	1.704, 1.014	0.769	0.860
Arg13	8.181	3.983	1.941, 1.735	1.606	3.162	7.189
Arg14	8.143	4.052	1.941, 1.733	1.725, 1.604	3.169	7.154
Ala15	8.422	4.114	1.500			
Ser16	8.110	4.275	4.029, 3.979			
Thr17	7.695	4.316	<u>4.373</u>	1.270		
Ile18	7.589	4.102	1.925	1.576, 1.216	0.851	0.912
Glu19	8.149	4.351	2.121, <u>2.011</u>	2.468, 2.426		
Met20	7.958	4.702	2.088	2.544		2.027
Pro21		4.487	2.539, 2.483	2.082, 1.948	3.876, 3.591	
Gln22	8.766	4.008	2.173	2.438		7.433, 6.742
Gln23	8.954	4.079	<u>2.084</u> , 1.945	2.464		7.457, 6.784
Ala24	7.455	4.155	1.474			
Arg25	7.858	3.944	1.876, 1.712	1.604	3.246, 3.181	7.253
Gln26	8.143	4.019	2.136, 2.115	2.461, 2.374		7.371, 6.721
Lys27	7.647	4.069	1.954, 1.598	1.613, 1.465	1.682	2.937, 7.604
Leu28	8.099	4.063	<u>1.773</u> , 1.712	1.510	0.849, 0.818	
Gln29	8.212	3.985	2.198, <u>2.117</u>	2.480, 2.367		7.172, 6.668
Asn30	7.908	4.482	<u>2.884</u> , 2.803		7.486, 6.821	
Leu31	7.878	4.154	1.718, <u>1.625</u>	1.557	0.859, 0.797	
Phe32	8.162	4.430	3.238, 3.096		7.173	7.197
Ile33	7.974	3.986	<u>1.930</u>	1.611, 1.283	0.900	0.891
Asn34	7.801	4.570	2.588, 2.511		7.337, 6.759	
Phe35	8.006	4.694	3.272, 3.058		7.277	7.207
Cys36	7.727	4.452	2.890			

TABLE VI. NMR assignments of cP226 (SEQ ID NO: 10) in water.

¹H-chemical shifts of cP226 (SEQ ID NO: 10) in 90% H₂O/10% D₂O, pH 6.50 at 10°C.

Values expressed in ppm.

res.	HN	HA	HB1*	HB2	HG*	HD1/ HD*	HD2/ HE*	HE3	HZ2
Cys 1		4.054	2.991	2.896					
Tyr 2	6.924	4.695	2.884			6.986	6.792		
Trp 3	8.094	4.599	3.199	3.087		7.106	10.024	7.427	7.465
Glu 4	8.313	3.907	1.909	1.785					
Leu 5	7.857	4.093	1.495	1.496	1.132	0.822			
Glu 6	8.227	4.123	2.032	1.982	2.195	7.074			
Trp 7	7.871	4.398	3.207			0.790	10.188	7.449	7.458
Leu 8	7.326	4.514	1.377		1.256				
Pro 9		4.179	2.214	1.820	1.974	3.617	3.505		
Cys 10	8.457	4.505	3.128	2.906					
Ala 11	8.062	4.115	1.330						

TABLE VII. Quality of the structure of cP226 (SEQ ID NO: 10) obtained by noe data

RMSD values were calculated from a family of 12 structures without NOE restraint violation bigger than 0.3 Å. The local RMSD was calculated on 3 residues

res.	Global displacem.			local RMSD	Local displacem.			NOEs (total)
	bb	heavy	heavysc		bb	heavy	heavysc	
Cys 1	1.80	1.99	1.77	0.00	0.00	0.00	0.00	2
Tyr 2	0.92	3.52	4.23	0.72	0.59	3.21	3.86	6
Trp 3	0.76	1.36	1.50	0.35	0.26	0.73	0.81	17
Glu4	0.83	2.22	2.86	0.29	0.23	1.77	2.33	12
Leu 5	0.63	1.38	1.78	0.26	0.20	1.04	1.44	23
Glu 6	0.73	2.34	3.00	0.26	0.24	1.73	2.23	9
Trp 7	0.65	2.06	2.39	0.27	0.25	1.67	1.96	16
Leu 8	0.53	1.55	2.11	0.18	0.16	1.58	2.22	27
Pro 9	0.60	0.83	1.04	0.17	0.12	0.18	0.21	15
Cys10	0.71	0.94	1.11	0.27	0.18	0.36	0.54	13
Ala11	1.20	1.67	1.73	0.00	0.00	0.00	0.00	6

Abbreviations

PLB, phospholamban; PLB[a.a.1-36], 36 a.a. N-terminal fragment of human phospholamban (SEQ ID NO:9); SR, sarcoplasmic reticulum; SERCA, sarco/endoplasmic reticulum Ca^{2+} -ATPase; SERCA_2, cardiac isoform of the sarco/endoplasmic reticulum Ca^{2+} -ATPase; CD, circular dichroism; COSY, correlation spectroscopy; TOCSY, total correlation spectroscopy; NOESY, nuclear Overhauser-enhancement spectroscopy; d_3 -TFE, perdeuterated trifluoroethanol; d_{10} -DTT, perdeuterated dithiotreitol.